

STATE OF ILLINOIS
HENRY HORNER, *Governor*
DEPARTMENT OF REGISTRATION AND EDUCATION

DIVISION OF THE
NATURAL HISTORY SURVEY

THEODORE H. FRISON, *Chief*

Vol. XX

BULLETIN

Article V

Annotated List of the Fishes of Illinois

D. JOHN O'DONNELL



PRINTED BY AUTHORITY OF THE STATE OF ILLINOIS

URBANA, ILLINOIS
AUGUST 1935

STATE OF ILLINOIS
HENRY HORNER, *Governor*
DEPARTMENT OF REGISTRATION AND EDUCATION
HONORABLE JOHN J. HALLIHAN, *Director*

BOARD OF NATURAL RESOURCES AND CONSERVATION
HONORABLE JOHN J. HALLIHAN, *Chairman*

WILLIAM TRELEASE, D.Sc., LL.D., *Biology*
HENRY C. COWLES, Ph.D., D.Sc., *Forestry*
JOHN W. ALVORD, C.E., *Engineering*
WILLIAM A. NOYES, Ph.D., LL.D., Chem.D.,
D.Sc., *Chemistry*
EDSON C. BASTIN, Ph.D., *Geology*
ARTHUR CUTTS WILLARD, D.Eng., LL.D.,
President of the University of Illinois

NATURAL HISTORY SURVEY DIVISION

URBANA, ILLINOIS

Scientific and Technical Staff

THEODORE H. FRISON, Ph.D., *Chief*

SECTION OF ECONOMIC ENTOMOLOGY

W. P. FLINT, B.S., *Chief Entomologist*
C. C. COMPTON, M.S., *Associate Entomologist*
M. D. FARRAR, Ph. D., *Research Entomologist*
S. C. CHANDLER, B.S., *Southern Field Entomologist*
J. H. BIGGER, B.S., *Central Field Entomologist*
L. H. SHROPSHIRE, M.S., *Northern Field Entomologist*
E. R. MCGOVAN, Ph.D., *Research Fellow in Entomology*
W. E. MCCAULEY, B. S., *Assistant Entomologist*
J. F. ALSTERLUND, M.S., *Research Fellow in Entomology*

SECTION OF APPLIED BOTANY AND
PLANT PATHOLOGY

L. R. TEHON, Ph.D., *Botanist*
J. C. CARTER, Ph.D., *Assistant Botanist*
G. H. BOEWE, B.S., *Field Botanist*

SECTION OF AQUATIC BIOLOGY

DAVID H. THOMPSON, Ph.D., *Zoologist*
F. D. HUNT, *Field Naturalist*
D. F. HANSEN, A.M., *Assistant Zoologist*
D. J. O'DONNELL, M.S., *Assistant Zoologist*

SECTION OF INSECT SURVEY

H. H. ROSS, Ph.D., *Systematic Entomologist*
CARL O. MOHR, Ph.D., *Associate Entomologist, Artist*
L. H. TOWNSEND, Ph.D., *Assistant Entomologist*

SECTION OF GAME RESEARCH AND
MANAGEMENT

R. E. YEATTER, Ph.D., *Game Specialist*

SECTION OF FORESTRY

L. E. SAWYER, B.S.F., *Extension Forester*


PUBLICATIONS

CARROLL CHOUINARD, M.A., *Editor*

This paper is a contribution from the Section of Aquatic Biology

CONTENTS

	PAGE
INTRODUCTION.....	473
CLASSIFIED LIST OF ILLINOIS FISHES.....	475
BIBLIOGRAPHY.....	491
APPENDIX.....	492
Hybridization and racial differentiation among Illinois fishes.—Index to families, genera and species.	



SURVEY LABORATORY BOAT ANAX ON THE ILLINOIS RIVER AT HAVANA

Annotated List of the Fishes of Illinois

D. JOHN O'DONNELL

SINCE 1876, specimens and data regarding the appearance and distribution of fishes in this state have been collected by zoologists of the ILLINOIS STATE NATURAL HISTORY SURVEY and the organizations which were its predecessors. In 1909 Stephen Alfred Forbes and Robert Earl Richardson published a report entitled "The Fishes of Illinois," which was partly revised for a second edition printed in 1920¹. Many scientific names of Illinois fishes have been changed in the twenty-six years that have passed since the original report was issued, so that today there is need for a list which will place the state fauna in accord with present general classifications.

Present Fauna

In recording the many changes that have been made by taxonomists, this list presents 172 species names, distributed among 31 families and 117 genera. In addition there are several dozen hybrid forms in the fish fauna of Illinois.

Ten species have been added to the forms known to Forbes and Richardson in 1909, some of which are names of American and foreign species that have become established in Illinois, and some native species previously undiscovered.

Jordan Nomenclature

Because of its general usage, the manual of Jordan (1929), which changed the generic names associated with 70 species and the specific names of 31 species of fishes given in Forbes & Richardson, has furnished the nomenclature for this list. Names in the list by Jordan, Evermann and Clark (1930), as well as the old names of Forbes & Richardson (1920), are used in synonymy.

Names of Hybrids

There are a few exceptions to this rule. One, of course, is the case of certain hybrids, which Jordan did not recognize but which have nevertheless been classified and used separately, either by Forbes and Richardson in 1909 or subsequently in Jordan, Evermann & Clark (1930). *Apomotis* (*Lepomis*) *ischyrus*, *A.* (*Lepomis*) *euryorus* and *Notropis piisbryi* are in this group. Likewise *Carpiodes thompsoni*, not recognized by Jordan and now probably best treated as a variety of *C. cyprinus*, is included in this list because it has been established in the literature of the Illinois fauna (Forbes & Richardson 1920), in addition to its placement as a distinct species in Jordan, Evermann & Clark (1930). *Alwordius* (*Hadropterus*) *evermanni* was included in Forbes & Richardson on the basis of a single

¹See bibliography, p. 491.



Fig. 1.—Interior of SURVEY laboratory boat Anax, looking forward.

specimen taken in the Illinois. Possibly a suspicion of its hybrid nature, as stated by Hubbs (1926), accounts for its omission from Jordan (1929). It recurs as valid, however, in Jordan, Evermann & Clark, on the strength of specimens from Indiana. The last exception is in the case of the red horses, whose names of Forbes & Richardson are retained. The descriptions of red horses given in Jordan (1929) and in Jordan, Evermann & Clark (1930), do not approximate our Illinois red horses closely enough to make their identity probable.

Forbes & Richardson Accurate

It has never been adequately emphasized that changes in the nomenclature of

Illinois fishes have resulted entirely from the formalities of fish taxonomy and not from correction of any mistakes in segregation or identifications as used in 1903-06 by Forbes and Richardson. Neither have they resulted from any evolution of our species, nor from any refinement in the method of describing them. The descriptions which were written for "The Fishes of Illinois" were, in almost every case, based on the detailed examination and measurement of large numbers of Illinois specimens.

Common Names in Illinois

Lack of agreement among ichthyologists in regard to common names of fishes has forced the adoption in this list of those

common names which are now in most general use in Illinois. These common names and the recorded distributions have been included as far as known; it is hoped that this effort will stimulate others interested in the fish life of the state to fill the obvious gaps in many of the records.

Acknowledgments

This report is based upon field records kept by SURVEY zoologists and their fore-runners, since 1876. A great deal of in-

formation and assistance have been contributed by present staff members, among them Dr. T. H. Frison, chief; Dr. David H. Thompson, zoologist; and Mr. F. D. Hunt, field naturalist, to all of whom the writer tenders thanks. An indebtedness is also acknowledged to Dr. Harley Jones Van Cleave, professor of zoology in the University of Illinois, for his suggestion of the problem while expressing a need among ichthyologists and less technical students for such a list.

CLASSIFIED LIST OF ILLINOIS FISHES

PETROMYZONIDAE

Ichthyomyzon Girard

I. concolor (Kirtland) *J JEC FR 9*²

LAMPER, LAMPER EEL, LAMPREY.—All Illinois streams with a drainage area exceeding 1000 square miles. Parasitic in winter on larger fishes such as carp, buffalo, catfishes and spoonbill. Three or more lampers may occasionally be found on one host fish.

Lethenteron Creaser & Hubbs

L. appendix (De Kay) *J JEC*

Lampetra wilderi Gage *FR 11*

BROOK LAMPREY, SMALL BLACK LAMPREY.—Very rare; Lake Michigan 1903, small creek near Danville winter of 1931. Numbers observed spawning on rocky riffles of Wild Cat creek near Dayton (Tippecanoe county), Ind., in early spring of 1910, 1916.

POLYODONTIDAE

Polyodon Lacépède

P. spathula (Walbaum) *J JEC FR 16*

SPOONBILL CAT, PADDLEFISH, BONELESS CAT.—Large rivers; recently found in the

Mississippi, Ohio, Wabash, Illinois river to Henry, Rock river to Sterling, Kaskaskia river to Evansville. Everywhere less abundant than formerly. Almost absent from the upper Mississippi following construction of Keokuk dam.

ACIPENSERIDAE

Acipenser Linnaeus

A. fulvescens Rafinesque *J JEC*

Acipenser rubicundus Le Sueur *FR 24*

ROCK STURGEON, LAKE STURGEON.—Taken occasionally in the Mississippi, Ohio, Wabash and Rock rivers. Common in the Illinois river before 1900; now rare.

Scaphirhynchus Heckel

S. platyrhynchus (Rafinesque) *J JEC FR 27*

SHOVEL-NOSED STURGEON, HACKLEBACK.—Common in the Mississippi and Ohio rivers. Occasionally taken in the lower Illinois.

Parascaphirhynchus Forbes & Richardson

P. albus Forbes & Richardson *J JEC FR 28*

WHITE STURGEON, LONG-NOSED STURGEON.—Rare; known only from the Mississippi river at Grafton and Alton.

²*J* indicates the present accepted name as given by Jordan (1929); *JEC* indicates the name given by Jordan, Evermann & Clark (1930); and *FR*, with a number, indicates the page on which the species is given in Forbes & Richardson (1920).



Fig. 2.—Principal streams of Illinois.

LEPISOSTEIDAE

Lepisosteus Lacépède

L. osseus (Linnaeus) *J JEC FR 31*

LONG-NOSED GAR, BILLY GAR.—Lakes. Found in all streams with drainage area exceeding 500 square miles.

Cylindrosteus Rafinesque

C. platostomus (Rafinesque) *J JEC*

Lepisosteus platostomus Rafinesque *FR 34*

SHORT-NOSED GAR.—Most abundant gar in rivers with drainage area of more than 5000 square miles.

Atractosteus Rafinesque

A. tristoechus (Schneider) *J*

Atractosteus spatula (Lacépède) *JEC*. *Lepisosteus tristoechus* (Bloch & Schneider) *FR 35*

ALLIGATOR GAR.—Mississippi river to Quincy. Occasionally in the Illinois to Henry, the Ohio, lower Wabash and lower Kaskaskia.

AMIIDAE

Amia Linnaeus

A. calva Linnaeus *J JEC FR 38*

DOGFISH, GRINDLE, BOWFIN.—Sloughs and lakes adjoining the Mississippi, Illinois and Ohio rivers. Not so abundant northward. Fox, Rock and Kankakee rivers. Occurs in the larger and more sluggish streams of southern Illinois.

HIODONTIDAE

Amphiodon Rafinesque

A. alosoides (Rafinesque) *J JEC*

Hiodon alosoides (Rafinesque) *FR 43*

NORTHERN MOONEYE, GOLDEYE.—Very rare; found only in the largest streams, and prefers swift open water. At one time abundant in the Mississippi and Ohio rivers, and the Illinois to Havana.

Hiodon Le Sueur

H. tergisus Le Sueur *J JEC FR 44*

TOOTHED HERRING, MOONEYE.—Occurs in small numbers in the Rock river

to Beloit, Illinois river to Henry, Kaskaskia and Wabash rivers. Very abundant in the Mississippi and Ohio.

CLUPEIDAE

Pomolobus Rafinesque

P. chrysochlorus Rafinesque *J JEC FR 48*

SKIPJACK, GOLDEN SHAD, BLUE HERRING.—Not common in Illinois, a few having been taken from the lower Rock and lower Illinois rivers. Present in small numbers in the Mississippi and Ohio.

Alosa Linck

A. ohiensis Evermann *J JEC FR 49*

OHIO SHAD.—Rare; present in small numbers in the Mississippi and the Ohio to Pittsburgh. May occur in the Illinois up to Meredosia.

DOROSOMIDAE

Dorosoma Rafinesque

D. cepedianum (Le Sueur) *J JEC FR 45*

GIZZARD SHAD, HICKORY SHAD.—Abundant; generally distributed in all larger rivers, connecting lakes and occasionally smaller streams. Abundant in lakes having mud bottom.

COREGONIDAE

Prosopium Milner

P. quadrilaterale (Richardson) *J JEC*

Coregonus quadrilateralis Richardson *FR 53*

MENOMINEE WHITEFISH, ROUND WHITEFISH.—Lake Michigan. Much more rarely taken than the common whitefish.

Coregonus Linnaeus

C. clupeaformis (Mitchill) *J JEC FR 51*

COMMON WHITEFISH, SUPERIOR WHITEFISH.—Fairly abundant in southwestern Lake Michigan. Largest and most widely known of our species of whitefish.

Leucichthys Dybowski**L. artedi** (Le Sueur) *J JEC**Argyrosomus artedi* (Le Sueur) *FR 54*

CISCO, LAKE HERRING, COMMON LAKE HERRING.—Lake Michigan. The most abundant food fish in the Great Lakes. In addition to the cisco, the following are more or less commonly taken in Lake Michigan: *L. reighardi* Koelz, the chub *L. zenithicus* (Jordan & Evermann), the longjaw *L. alpenae* Koelz, the mooneye cisco *L. hoyi* (Gill), *L. johanna* (Wagner), the mooneye *L. kiyi* Koelz, and the blackfin *L. nigripinnis* (Gill).

SALMONIDAE**Salmo** Linnaeus**S. salar sebago** Girard *J JEC*

LANDLOCKED SALMON, SEBAGO SALMON.—Lake Michigan: found in small numbers at the northern end; may also occur off Chicago.

Trutta Linnaeus**T. trutta** (Linnaeus) *J JEC*

GERMAN BROWN TROUT.—A few have been reported by anglers from the tributaries of the Kishwaukee and Pecatonica rivers. It is uncertain whether this trout is able to propagate itself in natural waters of Illinois.

T. iridea (Gibbons) *J**Salmo irideus* Gibbons *JEC*

RAINBOW TROUT.—Common off Indiana and Michigan shores of Lake Michigan. To be expected off the Illinois shore. Probably not separable from *Salmo gairdnerii* Richardson.

Cristivomer Gill & Jordan**C. namaycush** (Walbaum) *J JEC FR 56*

LAKE TROUT.—Fairly abundant in Lake Michigan off the Illinois shore.

Salvelinus Richardson**S. fontinalis** (Mitchill) *J JEC*

EASTERN BROOK TROUT, SPECKLED TROUT.—In Franklin spring seven miles northwest of Rockford, and in spring-fed

tributaries of the Kishwaukee, Pecatonica and Apple rivers. Is able to propagate itself in these localities. Also found in Blackberry creek near Yorkville but not propagating.

ANGUILLIDAE**Anguilla** Thunberg**A. bostoniensis** Le Sueur *J*

Anguilla rostrata (Le Sueur) *JEC*. *Anguilla chrysypa* Rafinesque *FR 59*

AMERICAN EEL, FRESH-WATER EEL.—

Found in all streams having a drainage basin exceeding 500 square miles. Up to several hundred pounds taken per day by commercial fishermen at Browning on the Illinois.

CATOSTOMIDAE**Cycleptus** Rafinesque**C. elongatus** (Le Sueur) *J JEC FR 65*

MISSOURI SUCKER, BLACK HORSE, BLUEFISH.—Mississippi, Ohio and lower Illinois rivers. Formerly common in the lower Rock.

Megastomatobus Fowler**M. cyprinella** (Cuvier & Valenciennes) *J JEC*

Ictiobus cyprinella (Cuvier & Valenciennes) *FR 68*

RED-MOUTH BUFFALO, BIG-MOUTH BUFFALO.—Very abundant in all larger streams and floodplain lakes. Found in the smaller, more sluggish streams with drainage areas exceeding 300 square miles. Second among Illinois commercial fishes.

Ictiobus Rafinesque**I. urus** (Agassiz) *J JEC FR 70*

MONGREL BUFFALO, ROUND BUFFALO.—Distribution same as big-mouth buffalo but it is less abundant. Present known facts concerning the life history of this fish permit the interpretation that it is a natural hybrid between *Megastomatobus cyprinella* and *I. bubalus*.

I. bubalus (Rafinesque) *J JEC FR 72*

SMALL-MOUTH BUFFALO, QUILLBACK

BUFFALO, HIGHBACK BUFFALO, RAZOR-BACK BUFFALO.—Common in the Mississippi, Ohio, Illinois and Rock, and in the principal streams of the state generally. Prefers deep water. Characteristically a channel fish although common in some bottomland lakes and sloughs. Commercially important.

Carpiodes Rafinesque

C. carpio (Rafinesque) *J JEC FR 76*

COMMON RIVER CARP, SHAD, SILVEREEN.—Common in the Illinois, Rock and Mississippi. Seldom ascends smaller streams.

C. difformis Cope *J JEC FR 77*

BLUNT-NOSED RIVER CARP.—Common in some smaller streams of central Illinois. Taken sparingly in the Rock and Illinois rivers.

C. cyprinus (Le Sueur) *J JEC*

Carpiodes velifer (Rafinesque) *FR 78*

QUILLBACK, SILVER CARP.—Common in medium-sized rivers and larger creeks. Most abundant in northern Illinois. Avoids area of the lower Illinoisan glaciation.

C. thompsoni Agassiz *JEC FR 79*

LAKE CARP.—Found in the Great Lakes. Its recognition by Forbes and Richardson from the Illinois river is questioned. A recent examination of this genus by the late Mr. R. E. Richardson and Dr. David H. Thompson, using a large amount of fresh material including specimens from several points on the Great Lakes and other parts of the United States, indicates that *C. thompsoni* should be regarded as a variety of *C. cyprinus*.

Catostomus Le Sueur

C. catostomus (Forster) *J JEC FR 84*

LONG-NOSED SUCKER, NORTHERN SUCKER, RED SUCKER.—Lower Lake Michigan.

C. commersonii (Lacépède) *J JEC FR 85*

BLACK SUCKER, FINE-SCALED SUCKER, COMMON SUCKER.—Generally distributed throughout Illinois. Most abundant in the northern third of the state, especially

in the smaller rivers and larger creeks. Common in Lake Michigan.

Hypentelium Rafinesque

H. nigricans (Le Sueur) *J JEC*

Catostomus nigricans Le Sueur *FR 86*

HOG SUCKER, HAMMERHEAD, STONE-ROLLER.—Common throughout northern and eastern parts of the state in headwaters of the smaller tributaries of the Illinois, Kaskaskia, Embarrass and Big Vermilion rivers. Prefers swifter streams with gravel bottom.

Erimyzon Jordan

E. sucetta oblongus (Mitchill) *J FR 81*

Erimyzon sucetta (Lacépède) *JEC*. *Erimyzon oblongus* (Mitchill) *JEC*

CHUB SUCKER, SWEET SUCKER.—Widely distributed in large and small streams. Essentially a creek species. Most abundant in eastern part of the state in the drainage of the Wabash and Ohio, and in the headwaters of the Sangamon and Kaskaskia. Common in the Fox lakes.

Minytrema Jordan

M. melanops (Rafinesque) *J JEC FR 83*

SPOTTED SUCKER, STRIPED SUCKER.—Taken in all stream systems, but most commonly in the Wabash and the Kaskaskia basins. Mainly a species of creeks and smaller rivers. Most abundant in certain lakes and ponds.

Moxostoma Rafinesque

M. anisurum (Rafinesque) *J JEC FR 89*

WHITE-NOSED SUCKER.—Lake Michigan and the Illinois, Sangamon, Rock, Mississippi, Ohio and Wabash. Distributed throughout the state in moderate numbers.

M. aureolum (Le Sueur) *J JEC FR 90*

COMMON RED HORSE.—Taken in all streams except the Big Muddy. Commonest in the northern and eastern two thirds of the state. Abundant in the Rock river basin and in the Kaskaskia, Wabash and Sangamon systems.

M. lesueurii (Richardson) *J JEC*

Moxostoma breviceps (Cope) *FR 91*

SHORT-HEADED RED HORSE.—All rivers, lakes and streams of the central and northern part. More abundant in the smaller rivers than in large rivers, creeks and lakes.

Placopharynx Cope

P. carinatus Cope *J JEC*

Placopharynx duquesnei (Le Sueur) *FR 93*

BIG-TOOTHED RED HORSE.—Rare in Illinois. One specimen from the Wabash, several from the Illinois. Not recognized with certainty in recent years.

^L
Hagochila Jordan & Brayton

^L
H. lacera Jordan & Brayton *FR 94*

RABBIT-MOUTH SUCKER.—Up to the present not found in Illinois. Probably inhabits the Wabash basin.

CYPRINIDAE

Cyprinus (Artedi) Linnaeus

C. carpio Linnaeus *J³ FR 104*

EUROPEAN CARP.—Common or abundant in all streams draining more than 200 square miles. It is very abundant in the Illinois river and its connecting bottomland lakes. The carp is the most important commercial fish of the Illinois, the annual catch being 6 to 10 million pounds.

Carassius Nilsson

C. auratus (Linnaeus) *J*

GOLDFISH.—Rarely taken from the Illinois river and Lake Michigan. Common in Wolf lake near Chicago. Three specimens of a hybrid *Carassius auratus* × *Cyprinus carpio* have been taken, two from the Illinois at Peoria in 1932 and one from the Rock at Rockford, Feb. 1, 1933.

Chrosomus Rafinesque

C. erythrogaster Rafinesque *J JEC FR 112*

RED-BELLIED DACE.—Locally abundant. Occurs in small, clear, spring-fed tributaries of the Rock and Fox rivers. Found

occasionally in small spring-fed streams of southern Illinois. Does not occur within the lower Illinoisan glaciation.

Notemigonus Rafinesque

N. crysoleucas (Mitchill) *J JEC*

Abramis crysoleucas (Mitchill) *FR 126*

GOLDEN SHINER, BREAM, ROACH.—Found in all streams. In the southern and eastern parts of the state it prefers creeks. Elsewhere it has been taken in greatest numbers along the larger rivers.

Opsopoeodus Hay

O. emiliae Hay *J JEC FR 124*

Most abundant in southern Illinois, especially in the Saline river. Prefers creeks and ponds. Northward it has been taken almost wholly along the larger rivers—the Illinois, the Rock and Mississippi.

Semotilus Rafinesque

S. atromaculatus (Mitchill) *J JEC FR 121*

HORNED DACE, CREEK CHUB.—Equally abundant in all the stream systems of Illinois. Prefers creeks and the smaller rivers.

Rhinichthys Agassiz

R. cataractae (Cuvier & Valenciennes) *J JEC FR 160*

LONG-NOSED DACE.—Rare; found only in Big creek near Anna, and near Waukegan. Prefers clear cold streams.

R. atronasus (Mitchill) *J JEC FR 162*

BLACK-NOSED DACE.—Rare; has been taken only in creeks of the northern part of the state, the Fox river, and from Big creek (two specimens) near Anna.

Extrarius Jordan

E. hyostomus (Gilbert) *J*

Macrhybopsis hyostoma (Gilbert) *JEC*

Hybopsis hyostomus (Gilbert) *FR 163*

Taken in moderate numbers from the Rock, Green, Illinois and Mississippi.

Macrhybopsis Cockerell & Alliston

M. gelidus (Girard) *J*

Macrhybopsis gelida (Girard) *JEC*

³See Jordan (1929), p. 67, footnote.

Newly discovered in Illinois. Several specimens were taken in February 1930 from the Mississippi river at Chester.

Erimystax Jordan

E. dissimilis (Kirtland) *J JEC*

Hybopsis dissimilis (Kirtland) *FR 164*

SPOTTED SHINER.—Rare; found in the lower Kaskaskia, upper Embarrass, the Sangamon, Kickapoo, Spoon and Rock.

Erinemus Jordan

E. hyalinus (Cope) *J JEC*

Hybopsis amblops (Rafinesque) *FR 165*

SILVER CHUB, BIG-EYED CHUB.—Abundant in the southeastern part, also in the Mackinaw and upper Kankakee rivers. Prefers creeks. Restricted in central Illinois to the Middle Fork, Salt Fork and Embarrass river basins.

E. storerianus (Kirtland) *J JEC*

Hybopsis storerianus (Kirtland) *FR 166*

STORER'S CHUB.—Scarce. Widely distributed in the larger streams and lowland lakes. Not present in streams of the lower Illinoisan glaciation.

Nocomis Girard

N. biguttatus (Kirtland) *J*

Nocomis kentuckiensis (Rafinesque) *JEC*.

Hybopsis kentuckiensis (Rafinesque) *FR 167*

RIVER CHUB, HORNYHEAD.—Found only in the recently glaciated areas. Very abundant in streams and lakes of the northeastern section. Hybridizes with *Luxilus cornutus*.

N. micropogon (Cope) *J JEC*

Has been taken only from streams of the Wabash drainage. This species may have been included with *N. biguttatus* by Forbes & Richardson under *Hybopsis kentuckiensis*.

Platygobio Gill

P. gracilis (Richardson) *J JEC FR 170*

FLATHEAD CHUB.—Has been taken only from the Ohio at Cairo and the Mississippi at Chester.

Hybopsis Agassiz

H. deliciosus (Girard) *J*

Hybopsis phenacobius Forbes *JEC*. *Notropis phenacobius* Forbes *FR 138*

Rare; streams of central Illinois.

H. volucellus (Cope) *J JEC*

Notropis blennius (Girard) *FR 137*

STRAW-COLORED MINNOW.—Distributed throughout in clear streams and creeks. Most abundant in the extreme northern part. Avoids the southern Illinoisan glaciation.

H. dorsalis (Agassiz) *J*

Hybopsis gilberti (Jordan & Meek) *JEC*.

Notropis gilberti Jordan & Meek *FR 139*

Prefers small rivers and creeks. Found in the Mississippi drainage.

H. boops Gilbert *J*

Hybopsis shumardi (Girard) *JEC*. *Notropis illecebrosus* (Girard) *FR 140*

BIG-EYED MINNOW.—Rare; closely limited to the tributaries of the Wabash in the eastern part.

H. anogenus (Forbes) *J JEC*

Notropis anogenus Forbes *FR 132*

PUGNOSE SHINER.—Rare in Illinois. Twenty-four specimens were collected in the upper Fox river at McHenry, Ill., in 1885, one individual from Fourth lake in 1892, three individuals from Pistakee lake in 1925 and 15 in 1930 from the same water.

H. heterolepis Eigenmann & Eigenmann *J*

Hybopsis cayuga (Meek) *JEC*. *Notropis cayuga* Meek *FR 133*

CAYUGA SHINER, BLACK-NOSED SHINER.—Not abundant in Illinois. Most common in creeks in the northern half of the state, although found in the north-eastern glacial lakes.

H. heterodon (Cope) *J JEC*

Notropis heterodon (Cope) *FR 134*

BLACK-NOSED SHINER.—Distributed sparingly throughout, mainly in the lowland and glacial lakes in a way to indicate an avoidance of the lower Illinoisan glaciation. There is some evidence that one or two varieties of this species occur in Illinois but at the present time we are not able to separate them geographically or taxonomically.

H. nux richardsoni Hubbs & Greene *J JEC*

This form was found by Forbes and Richardson but the species had not been named at the time of their final report. It is closely related to *H. heterodon*. Taken in the lowland streams of the Wabash, Ohio and Big Muddy valleys.

H. atrocaudalis (Evermann) *J JEC*

Notropis cayuga atrocaudalis Evermann *FR 134*

Ten collections from the Illinois and adjacent waters near Meredosia, and one from the main river at Havana. Found in small numbers in the Kaskaskia and the streams of Champaign county.

H. hudsonius (DeWitt Clinton) *J*

Hudsonius hudsonius (DeWitt Clinton) *JEC*.

Notropis hudsonius (DeWitt Clinton) *FR 141*

SPOT-TAILED MINNOW.—Very abundant though limited to the Mississippi and Lake Michigan drainage, and occurring but twice south of the central part of the state. Most abundant in the larger rivers and in lakes.

H. blennius (Girard) *J*

Paranotropis jejunus (Forbes) *JEC*. *Notropis jejunus* (Forbes) *FR 150*

Distributed sparingly along the larger streams. Scarce in creeks and only moderately abundant in lowland lakes. Avoids the lower Illinoisan glaciation.

Cyprinella Girard

C. lutrensis (Baird & Girard) *J*

Moniana lutrensis (Baird & Girard) *JEC*.

Notropis lutrensis (Baird & Girard) *FR 143*

REDFIN.—Essentially a western species, occurring only in streams of the Mississippi drainage. Prefers streams of all sizes to lakes. Forbes & Richardson (1920) states, "It is very nearly allied to the next species, *N. whipplii*, compared with which it seems to be merely a more specialized form, the two sometimes intergrading in an obscure and very puzzling way."

C. whipplii (Girard) *J*

Erogala whipplii (Girard) *JEC*. *Notropis whipplii* (Girard) *FR 145*

STEEL-COLORED MINNOW.—Extremely abundant, especially in the smaller streams of the central part. Occurs in small numbers in all other parts. Scarce in lakes and large rivers. Prefers swift water.

Luxilus Rafinesque

L. cornutus (Mitchell) *J JEC*

Notropis cornutus (Mitchill) *FR 147*

COMMON SHINER.—Very abundant. Prefers creeks and the smaller rivers. Scarce in the southern third of the state, abundant in the rest. Hybridizes with *Nocomis biguttatus*, *Hybognathus nuchalis* and *Notropis rubrifrons*.

Notropis Rafinesque

N. photogenis (Cope) *J JEC*

SILVER SHINER.—Recently found in bottomland lakes of the Ohio near Brookport.

N. rubrifrons (Cope) *J JEC FR 153*

ROSY-FACED SHINER.—Occurs only in the Mississippi drainage of the northern third of the state. Taken in the tributaries of the Illinois, the Rock, the Fox and the Mississippi. Prefers clear rapid water. Hybridizes with *Luxilus cornutus*.

N. atherinoides Rafinesque *J JEC FR 151*

LAKE SHINER.—Distributed throughout. Less abundant in the Illinois, the Kaskaskia and the Big Muddy than in the other stream systems. Not present in the glacial lakes of northeastern Illinois.

N. pilsbryi Fowler *JEC FR 149*

This species is a hybrid between *Luxilus cornutus* and *Notropis rubrifrons*. Taken only in the east fork of the Mazon river near Gardner, and from the Sangamon in Champaign county.

Lythrurus Jordan

L. umbratilis cyanocephalus (Copeland) *J*

Lythrurus umbratilis (Girard) *JEC*. *Lythrurus atripes* Jordan *JEC*. *Notropis umbratilis atripes* (Jordan) *FR 154*

BLACKFIN.—Distributed throughout, being several times more numerous in the eastern half than in the western. Most

abundant in the Big Muddy, the tributaries of the Wabash and the small rivers and creeks of extreme southern Illinois. Jordan (1929) described this species as *Lythrurus atripes* from collections taken in streams of Union and Johnson counties.

Ericymba Cope

E. buccata Cope *J JEC FR 156*

SILVER-MOUTH MINNOW.—Headwaters of the minor tributaries of the Wabash and upper course of the Kaskaskia. Also in the tributaries of the Iroquois and the upper Sangamon.

Phenacobius Cope

P. mirabilis (Girard) *J JEC FR 158*

SUCKER-MOUTH MINNOW.—Most abundant in creeks. Occurs in all of our river basins but is not present in the upland glacial lakes. Prefers swift water and sandy bottom.

Ceraticthys Baird & Girard

C. vigilax Baird & Girard *J JEC*

Cliola vigilax (Baird & Girard) *FR 128*

BULLHEAD MINNOW.—Common throughout. Most abundant in the Kaskaskia and Wabash basins. Prefers swifter streams with clean bottom.

Hyborhynchus (Agassiz)

H. notatus (Rafinesque) *J JEC*

Pimephales notatus (Rafinesque) *FR 119*

BLUNT-NOSED MINNOW.—The most abundant and widely distributed minnow in Illinois. Very abundant in the lower Illinoisan glaciation, slightly less so in the waters of the northeastern part of the state.

Pimephales Rafinesque

P. promelas Rafinesque *J JEC FR 117*

BLACKHEAD MINNOW, FATHEAD.—Found in tributaries of all the stream systems in the northern and western three fourths of the state. Most abundant in creeks, the smaller rivers and ponds.

Hybognathus Agassiz

H. nuchalis Agassiz *J JEC FR 114*

SILVERY MINNOW.—Generally distributed throughout, occurring in many stream systems but most abundantly in southern Illinois. Hybridizes with *Luxilus cornutus*.

Dionda Girard

D. nubila (Forbes) *J JEC*

Hybognathus nubila (Forbes) *FR 116*

Rare; taken only in the northern part—Rock and Fox rivers.

Campostoma Agassiz

C. anomalum (Rafinesque) *J JEC FR 110*

DOUGHBELLY, GREASER CHUB, STONE-ROLLER.—Found in all creeks, where it is abundant, in the smaller rivers and occasionally in the larger rivers and lakes. Not found in streams of the lower Illinoisan glaciation.

AMEIURIDAE

Ictalurus Rafinesque

I. furcatus (Cuvier & Valenciennes) *J JEC FR 178*

BLUE CAT, FULTON CAT.—Common in the Mississippi and Ohio rivers and for a distance up their larger tributaries.

I. anguilla Evermann & Kendall *J JEC FR 179*

EEL CAT, WILLOW CAT.—Common. Found with *I. punctatus* and *Opladelus olivaris*, but in smaller numbers throughout the range.

I. punctatus (Rafinesque) *J JEC FR 180*

CHANNEL CAT, FIDDLER, SILVER CAT.—The most abundant of our true catfishes. It occurs in all river basins, sloughs and lakes. Four specimens taken in the Rock river at Rockford and Sterling in recent years seem to be hybrids between *Ictalurus punctatus* and *Opladelus olivaris*.

Villarius Rutter

V. lacustris (Walbaum) *J*

Haustor lacustris (Walbaum) *JEC. Ameiurus lacustris* (Walbaum) *FR 184*

CATFISH OF THE LAKES, NORTHERN CATFISH.—Reported as common in Lake Michigan.

Ameiurus Rafinesque

A. natalis (Le Sueur) *J JEC FR 187*

YELLOW BULLHEAD, YELLOW CAT.—Abundant throughout, except in the extreme northwestern part of the state. Most common in creeks and lowland lakes but found in all river basins, including the Michigan drainage area and the north-eastern glacial lakes.

A. nebulosus (Le Sueur) *J JEC FR 187*

SPECKLED BULLHEAD, BROWN BULLHEAD.—Distributed throughout in lakes and large sluggish rivers, except in the extreme northwestern part of the state and within the lower Illinoisan glaciation. The least abundant of our bullheads.

A. melas (Rafinesque) *J JEC FR 190*

BLACK BULLHEAD.—Distributed throughout, very abundant in the smaller streams. Least abundant in the streams of the Lake Michigan drainage and most abundant in the creeks of the Mississippi bluffs and in the valleys of the Wabash and Kaskaskia rivers. The most abundant of our bullheads.

Opladelus Rafinesque

O. olivaris (Rafinesque) *J JEC*

Leptops olivaris (Rafinesque) *FR 193*

MUD CAT, FLATHEAD CAT, GOUJON, YELLOW CAT.—Common in the Illinois, Mississippi and Ohio, and also found in the Rock and Wabash rivers. Taken in smaller numbers from the Spoon, Green, Little Wabash, Sangamon and Kaskaskia rivers and from Crooked creek.

Noturus Rafinesque

N. flavus Rafinesque *J JEC FR 194*

STONECAT.—Common and widely distributed in the larger creeks and smaller rivers of the northern half of the state, but has never been taken south of Douglas county. Prefers swift current and a rock and gravel bottom.

Rabida Jordan

R. exilis (Nelson) *J JEC*

Schilbeodes exilis (Nelson) *FR 199*

SLENDER STONECAT.—Rare; found in

the Illinois river, in the Pecatonica at Freeport, in the Du Page in Will county, in Honey creek in Henderson county, in the Sangamon in Champaign county, and in two creeks of Union county in extreme southern Illinois.

R. miurus (Jordan) *J JEC*

Schilbeodes miurus (Jordan) *FR 200*

BRINDLED STONECAT.—Confined principally to tributaries of the Wabash and Ohio. Although common, it has a spotty distribution in streams, preferring a gravel bottom and swift current. It is occasionally found in the extreme headwaters of the Kaskaskia.

Schilbeodes Bleeker

S. gyrinus (Mitchill) *J JEC FR 197*

TADPOLE CAT, MAD TOM.—Distributed throughout in larger rivers, creeks, upland and lowland lakes. Prefers still and muddy water. Enters the lower Illinoisan glaciation. This species is most abundant south and east in branches of the Kaskaskia and Wabash rivers.

S. nocturnus (Jordan & Gilbert) *J FR 198*

Rabida nocturna (Jordan & Gilbert) *JEC*

FRECKLED STONECAT.—Rare; found in creeks near Havana and Lincoln, in tributaries of the Kaskaskia in Clinton and Shelby counties, and in Camp creek in Henderson county.

UMBRIDAE**Umbra** (Krämer) Müller

U. limi (Kirtland) *J JEC FR 203*

MUD MINNOW, MUDFISH.—Occurs almost entirely in ponds, lakes and ditches in the extreme northern and southern parts. Occasionally taken at Havana and Meredosia on the Illinois river.

ESOCIDAE**Esox** (Artedi) Linnaeus

E. vermiculatus Le Sueur *J JEC FR 206*

GRASS PIKE, LITTLE PICKEREL.—Dis-

tributed throughout, most abundantly in the southern part. Most common in creeks, but also found in ponds and the smaller rivers. Prefers quiet and muddy waters.

E. lucius Linnaeus *J FR 207*

COMMON PIKE, PICKEREL, GREAT NORTHERN PIKE.—Abundant in the headwaters of the Kankakee and the glacial lakes of the northeastern part. Occasionally taken in the clearer sloughs and lakes of the Illinois, Rock and Green rivers. This may prove to be *Esox estor* Le Sueur, an American form as yet not clearly separated from *Esox lucius* Linnaeus of Europe, as is stated in Jordan, Evermann & Clark (1930).

E. masquinongy Mitchill *J JEC FR 209*

MUSKELLUNGE.—Reported in earlier years from lakes in the northeastern part. One was taken from Fox lake by a SURVEY field party in 1930.

CYPRINODONTIDAE

Fundulus Lacépède

F. diaphanus menona (Jordan & Copeland) *J FR 211*

Zygionectes diaphanus menona (Jordan & Copeland) *JEC*

MENONA TOP MINNOW, MENONA KILLIFISH.—A northern and middle western species in the United States. Has been taken here less than 50 times, all in the northern half of the state. Found in the headwaters of the Fox, Des Plaines and Rock rivers, in pools near Bloomington and in the Calumet ponds. Prefers clear water and sand bottom.

Zygionectes Agassiz

Z. dispar Agassiz *J JEC*

Fundulus dispar Agassiz *FR 212*

BLACKCHEEK TOP MINNOW.—Occurs throughout in bottomland lakes of the larger rivers and in creeks. Abundant in the upland lakes of northeastern Illinois. Most abundant in lakes and sloughs.

Z. notatus (Rafinesque) *J JEC*

Fundulus notatus (Rafinesque) *FR 213*

TOP MINNOW, TOPWATER.—Very abundant throughout in all types of waters. Most abundant in the smaller streams and headwaters of southern and eastern parts.

POECILIIDAE

Gambusia Poey

G. patruelis (Baird & Girard) *J JEC*

Gambusia affinis (Baird & Girard) *FR 215*

MOSQUITO FISH, VIVIPAROUS TOP MINNOW.—Fairly common in extreme southern Illinois. It has been taken also from Quincy, Meredosia, Pekin and in the Fox river basin. Several specimens taken from Horseshoe lake, July 5, 1933, contained embryos with prominent eye spots. Females collected August 15 contained young apparently ready for extrusion. Prefers ditches, marshes and lagoons.

AMBLIOPSIDAE

Forbesella Jordan & Evermann

"This genus contains transitional species connecting *Chologaster* with *Typhlichthys*," Jordan (1929).

F. papillifera (Forbes) *J JEC*

Chologaster papilliferus Forbes *FR 218*

SPRING CAVE FISH.—Cave springs, Union and Pope counties.

PERCOPSIDAE

Percopsis Agassiz

P. omiscomaycus (Walbaum) *J JEC*

Percopsis guttatus Agassiz *FR 225*

TROUT PERCH.—Abundant in bottomland lakes along the Illinois river from Meredosia to Hennepin. Also from a small stream near Lincoln, Logan county, and once from Lake Michigan off Chicago.

APIREDODERIDAE

Aphredoderus Le Sueur

A. sayanus (Gilliams) *J JEC FR 229*

PIRATE PERCH.—In muddy pools and streams throughout the state, most com-

mon southward. Prefers muddy bottom and little or no current.

CASTEROSTEIDAE

Pungitius Coste

P. pungitius (Linnaeus) *J JEC*

Pygosteus pungitius (Linnaeus) *FR 224*

NINE-SPINE STICKLEBACK.—Taken but once, from the lower Calumet river and from the adjacent part of Lake Michigan.

Eucalia Jordan

E. inconstans (Kirtland) *J JEC FR 222*

BROOK STICKLEBACK.—Confined to the lakes of northeastern Illinois, the Calumet river at South Chicago, tributaries of the Rock and Fox rivers, and creeks in La Salle county. Prefers clear cool brooks. Associated with water cress.

ATHERINIDAE

Labidesthes Cope

L. sicculus (Cope) *J JEC FR 227*

BROOK SILVERSIDE.—Distributed throughout the northern, central and eastern parts, but not taken from the Big Muddy, the Saline and other waters of extreme southern Illinois. Prefers clear streams and lakes.

CENTRARCHIDAE

Huro Cuvier

H. salmoides (Lacépède) *J*

Huro floridana (Le Sueur) *JEC. Micropterus salmoides* (Lacépède) *FR 267*

LARGE-MOUTH BLACK BASS.—Equally distributed throughout, passing freely into the lower Illinoisan glaciation. Most common in the larger and more sluggish rivers, upland and lowland lakes.

Micropterus Lacépède

M. dolomieu Lacépède *J JEC FR 263*

SMALL-MOUTH BLACK BASS.—A northern fish, avoiding the lower Illinoisan glaciation. Most abundant in the smaller rivers and about half as abundant in creeks. Prefers clear swift water.

M. pseudaplites Hubbs *J JEC*

KENTUCKY BASS, SOUTHERN SMALL-MOUTH BLACK BASS.—Two specimens taken April 16, 1935, from Grand Pierre creek near Rosiclaire and one from the Ohio river at Shawneetown. General distribution is southeastern United States from western West Virginia northward to southern Ohio, westward to Kansas and central Texas.

Apomotis Rafinesque

A. cyanellus ((Rafinesque) *J JEC*

Lepomis cyanellus Rafinesque *FR 248*

GREEN SUNFISH, BLUE-SPOTTED SUNFISH.—Common throughout. Most abundant in creeks, the smaller rivers and ponds. Hybridizes with *Eupomotis gibbosus*, *Allotis humilis*, *Helioperca incisor*, *Xenotis megalotis* and *Chaenobryttus gulosus*.

A. ischyryus (Jordan & Nelson) *JEC*

Lepomis ischyryus (Jordan & Nelson) *FR 250*

BIG-NOSED SUNFISH.—Rare; not known outside of Illinois. Several specimens have been taken from the Illinois river at Meredosia. Proved by Hubbs (1932) and recent SURVEY experiments to be a hybrid between *Apomotis cyanellus* and *Helioperca incisor*.

A. euryorus (McKay) *JEC*

Lepomis euryorus McKay *FR 252*

Very rare. Has been taken only from Crooked creek near La Harpe, in Hancock county. Hubbs (1932) has proved by experimental breeding that this form is the hybrid *Apomotis cyanellus* × *Eupomotis gibbosus*.

Lethogrammus Hubbs

L. symmetricus (Forbes) *J*

Apomotis symmetricus (Forbes) *JEC. Lepomis symmetricus* Forbes *FR 251*

Rare; found in lakes and streams of southern Illinois. Also taken twice from the Illinois river at Pekin.

Sclerotis Hubbs

S. miniatus (Jordan) *J JEC*

Lepomis miniatus Jordan FR 253

OLD-FASHIONED SUNFISH, RED PERCH.—Formerly common, now rare; taken occasionally from bottomland lakes and ponds of the Illinois river.

Allotis Hubbs

A. humilis (Girard) J JEC

Lepomis humilis (Girard) FR 255

ORANGE-SPOTTED SUNFISH.—Distributed throughout, most abundant in creeks and the smaller rivers. Rare or absent in extreme northern Illinois. Hybridizes with *Apomotis cyanellus*, *Helioperca incisor* and *Xenotis megalotis*. In Lake Senachwine and in Crystal lake in Urbana, it shows intergrades of many sorts with *Eupomotis gibbosus*, as if some hybrids between these species at these points had proved fertile and produced other segregates.

Xenotis Jordan

X. megalotis (Rafinesque) J JEC

Lepomis megalotis (Rafinesque) FR 254

LONG-EARED SUNFISH.—Extremely abundant in the smaller streams of the southern and eastern parts, comparatively scarce in the remainder of the state. Hybridizes with *Eupomotis gibbosus*, *Helioperca incisor*, *Apomotis cyanellus* and *Allotis humilis*.

X. megalotis peltastes (Cope) J

Lepomis megalotis peltastes Cope FR 254

DWARF LONG-EARED SUNFISH.—A dwarf race of northern Illinois. Has been taken only in the clear swift water of the Fox from Ottawa to the Fox lakes, in the Du Page at Naperville, in the Vermilion at Pontiac and Fairbury, in a small creek in Du Page county and in Indian creek, La Salle county.

Helioperca Jordan

H. incisor (Cuvier & Valenciennes) J JEC

Lepomis pallidus (Mitchill) FR 257

BLUEGILL, BLUE SUNFISH.—Principal sunfish of the state. Occurs throughout but most abundant in the larger streams and their principal tributaries. Also com-

mon in all lakes. Hybridizes with *Eupomotis gibbosus*, *Xenotis megalotis*, *Apomotis cyanellus*, *Chaenobryttus gulosus* and *Allotis humilis*.

Eupomotis Gill & Jordan

E. heros (Baird & Girard) J JEC FR 259

A southern species, rare in Illinois. Has been taken at a few points in the Wabash basin and in the Ohio at Brookport.

E. gibbosus (Linnaeus) J JEC FR 260

PUMKINSEED.—Very abundant in the upland lakes of Lake and McHenry counties and in lakes along the Illinois. Very scarce south of the center of the state. Essentially a pond species, and next most abundant in the smaller rivers. Produces sterile hybrids with *Helioperca incisor*, *Xenotis megalotis*, *Apomotis cyanellus* and *Chaenobryttus gulosus*. Shows fertile segregates with *Allotis humilis*.

Chaenobryttus Gill

C. gulosus (Cuvier & Valenciennes) J JEC FR 245

WARMOUTH BASS.—Distributed throughout in small numbers. Abundant in the glacial lakes of northeastern Illinois and also in the lower Illinoisan glaciation. Prefers muddy water. Hybridizes with *Eupomotis gibbosus*, *Helioperca incisor* and *Apomotis cyanellus*.

Ambloplites Rafinesque

A. rupestris (Rafinesque) J JEC FR 243

ROCK BASS, GOGGLE-EYE.—Mainly a northern species, having been taken from only five localities in southern Illinois, and not at all in the lower Illinoisan glaciation. Most abundant in rivers of medium size and in the glacial lakes.

Pomoxis Rafinesque

P. annularis Rafinesque J JEC FR 238

WHITE CRAPPIE.—Occurs in all parts, most abundantly in lakes, ponds and bayous but common also in the smaller rivers and in creeks.

P. sparoides (Lacépède) J JEC FR 240

BLACK CRAPPIE, CALICO BASS.—Found

throughout, usually with the white crap-
pie. It is slightly less common than *P.*
annularis in creeks, but more abundant in
the northeastern glacial lakes and the bot-
tomland lakes of the Illinois.

Centrarchus Cuvier & Valenciennes

C. macropterus (Lacépède) *J JEC FR 241*

ROUND SUNFISH, FLIER.—Found only
in southern Illinois from Hamilton coun-
ty southward; taken only from creeks and
sloughs tributary to the Little Wabash,
the Big Muddy, the Cache and a small
creek near Brookport. Very abundant in
Horseshoe lake.

ELASSOMIDAE

Elassoma Jordan

E. zonatum Jordan *J JEC FR 232*

PIGMY SUNFISH.—Has been taken only
six times; from Little Fox river at Phil-
lipstown, Wabash river at Wabash Sta-
tion, Drew pond in White county, Swan
pond near St. Francisville, Running lake,
and from a bluff spring in Union county.

ETHEOSTOMIDAE

Vigil Jordan

V. pellucidus (Baird) *J JEC*

Ammocrypta pellucida (Baird) *FR 301*

SAND DARTER.—Has been taken only in
the Middle Fork of the Vermilion in the
Wabash basin, lower Kaskaskia, lower
Illinois and in the Rock river.

Crystallaria Jordan & Gilbert

C. asprella (Jordan) *J JEC FR 300*

Rare; has been taken from the Rock
river at Cleveland, Erie and Milan, from
the Little Wabash at Effingham and the
Mississippi at East Dubuque. First de-
scribed from specimens taken in a rocky
creek of the Mississippi bluffs in Hancock
county.

Boleosoma De Kay

B. nigrum (Rafinesque) *J JEC FR 294*

JOHNNY DARTER.—Very abundant, dis-

tributed throughout. Most abundant in
the small streams of the Kaskaskia and
Wabash systems. Prefers creeks and
small brooks.

Vaillantia Jordan

V. camura (Forbes) *J JEC*

Boleosoma camurum Forbes *FR 298*

SNUB-NOSED DARTER.—Very abundant
in southern part. Most abundant in the
Big Muddy and Saline basins. Prefers
creeks and the smaller rivers.

Etheostoma Rafinesque

E. blennioides Rafinesque *J JEC*

Diplesion blennioides (Rafinesque) *FR 292*

GREEN-SIDED DARTER.—Has been taken
only from the smaller streams of the
Wabash system.

Imostoma Jordan

I. shumardi (Girard) *J JEC*

Cottogaster shumardi (Girard) *FR 290*

RIVER DARTER.—Not common. Occurs
wholly along the courses of the larger
streams, and has been taken from Lake
Michigan at Chicago.

Percina Haldeman

P. caprodes (Rafinesque) *J JEC FR 282*

LOG PERCH.—Distributed throughout.
Most common in lakes and medium-sized
rivers. More abundant in the northern
than in central and southern parts.

Alvordius Girard

A. evermanni (Moenkhaus) *JEC*

Hadropterus evermanni Moenkhaus *FR 284*

One specimen taken from the Illinois
river at Havana. Hubbs (1926) regards
this form as a hybrid of undetermined
parentage.

A. phoxocephalus (Nelson) *J JEC*

Hadropterus phoxocephalus (Nelson) *FR 285*

SHARP-NOSED DARTER.—Occurs
throughout except in the northern glacial
lakes. Most abundant in the smaller
streams.

A. maculatus Girard *J JEC*

Hadropterus aspro (Cope & Jordan) *FR 286*

BLACK-SIDED DARTER.—Occurs

throughout except in the northern glacial lakes. Common in the smaller rivers and creeks, rare in the larger rivers.

A. ouachitae (Jordan & Gilbert) *J JEC*
Hadropterus ouachitae (Jordan & Gilbert)
FR 288

One specimen taken from the Wabash river at New Harmony, Indiana, in 1909.

Serraria Gilbert

S. sciera (Swain) *J JEC*
Hadropterus scierus Swain *FR 289*

Two specimens taken from the Embarrass river near Charleston.

Ericosma Jordan

E. evides (Jordan & Copeland) *J JEC*
Hadropterus evides (Jordan & Copeland)
FR 288

BARRED DARTER.—One specimen taken from the Rock river in 1877. Several specimens taken from the same river at Rock Island in 1927 and at Sterling in 1932. One specimen taken from the Kaskaskia in 1930.

Nanostoma Putnam

N. zonale (Cope) *J JEC*
Etheostoma zonale (Cope) *FR 304*

BANDED DARTER.—Limited to the northern half of the state except for a single collection from the Wabash. Prefers the smaller rivers to creeks or large rivers.

Nothonotus Agassiz

N. camurus (Cope) *J JEC*
Etheostoma camurum (Cope) *FR 306*

BLUE-BREASTED DARTER.—Has been taken several times from the Salt Fork and Middle Fork of the Vermilion river. Reported from Peoria, from Union county and the Saline and lower Wabash basins.

Oligocephalus Girard

O. jessiae (Jordan & Brayton) *J JEC*
Etheostoma jessiae (Jordan & Brayton) *FR 307*

MUD DARTER.—Abundant. Prefers large streams and lowland lakes to small rivers and creeks. Not found in the up-

land glacial lakes. Three times as abundant in central and nearly twice as abundant in southern, as in northern Illinois.

O. coeruleus (Storer) *J JEC*
Etheostoma coeruleum Storer *FR 309*

RAINBOW DARTER.—Prefers creeks and smaller rivers having swift water and a clean bottom. More abundant in northern Illinois than is *O. jessiae*. Avoids the lower Illinoisan glaciation.

O. obeyensis (Kirsch) *J*
Nivicola obeyense (Kirsch) *JEC. Etheostoma obeyense* Kirsch *FR 311*

Rare; has been taken only in four collections, all from rocky and gravelly creeks in Pope and Hardin counties.

O. exilis (Girard) *J*
Oligocephalus iowae (Jordan & Meek) *JEC. Etheostoma iowae* Jordan & Meek *FR 306*

Abundant in northern Illinois in Pistakee lake, Wolf lake, Senachwine lake, Rock river, Green river, Pecunsagan creek, in the Illinois river at Ottawa and Dutchman's creek in Johnson county.

Claricola Jordan & Evermann

C. squamiceps (Jordan) *J JEC*
Etheostoma squamiceps Jordan *FR 312*

Has been taken only in southern Illinois south of the Saline river, except for one collection from Robinson creek, a branch of the Kaskaskia in Shelby county, and a collection from the Little Wabash river near Carmi in White county. Prefers swift water and gravel bottom.

Catonotus Agassiz

C. flabellaris (Rafinesque) *J JEC*
Etheostoma flabellare Rafinesque *FR 313*

FAN-TAILED DARTER.—Prefers the smaller streams. Has been taken only once within the lower Illinoisan glaciation. Most abundant in the northern third of the state, fairly abundant in the central third.

Hololepis Agassiz

H. fusiformis (Girard) *J*
Boleichthys fusiformis (Girard) *JEC FR 315*

FUSIFORM DARTER, BROWN-SIDED DART-

ER.—Most abundant in the creeks and smaller rivers. Found in the upland lakes of Lake and McHenry counties and abundant in the southeastern part of the state. Prefers sluggish water and mud bottom.

Microperca Putnam

M. punctulata Putnam *J JEC FR 317*

LEAST DARTER.—Not present in the central part but abundant in the upland lakes of northeastern Illinois. Has been taken from Skillet Fork in Wayne county and from Drury creek in Union county.

PERCIDAE

Perca (Artedi) Linnaeus

P. flavescens (Mitchill) *J JEC FR 276*

YELLOW PERCH, RINGED PERCH.—Common in the upland lakes of the northeastern part and in the Illinois and Mississippi rivers as far south as Meredosia. Has never been taken in the Wabash or Kaskaskia systems or from streams in southern Illinois.

Stizostedion Rafinesque

S. vitreum (Mitchill) *J JEC FR 272*

WALL-EYED PIKE, WALLEYE, JACK SALMON.—Abundant in the swifter waters of the Rock and Kankakee rivers. Smaller numbers in the Wabash, Mississippi, Fox and Illinois rivers.

Cynoperca Gill & Jordan

C. canadensis grisea (De Kay) *J*

Cynoperca grisea (De Kay) *JEC. Stizostedion canadense griseum* (De Kay) *FR 274*

GRAY PIKE, SAUGER, SAND PIKE.—Has been taken mainly from the Mississippi and Illinois and sparingly from the Rock, Wabash and Kaskaskia. Most common in the Ohio. In this state, *C. grisea* probably grades into *C. canadense*.

MORONIDAE

Lepibema Rafinesque

L. chrysops (Rafinesque) *J JEC*

Roccus chrysops (Rafinesque) *FR 319*

WHITE BASS.—Larger rivers and lakes. Most abundant in the Fox lakes and the Mississippi, Illinois and Ohio rivers.

Chrysoperca Fowler

C. interrupta (Gill) *J JEC*

Morone interrupta Gill *FR 321*

YELLOW BASS.—Prefers the large rivers and adjacent lakes. Most abundant in central Illinois.

SCIAENIDAE

Aplodinotus Rafinesque

A. grunniens (Rafinesque) *J JEC FR 323*

SHEEPSHEAD, FRESH-WATER DRUM, WHITE PERCH.—Abundant. Distributed throughout, commonest in central part.

COTTIDAE

Cottus (Artedi) Linnaeus

C. bairdii Girard *J*

Cottus ictalops (Rafinesque) *JEC FR 326*

MILLER'S THUMB, COMMON SCULPIN.—Has been taken only in northern and southern Illinois: in McHenry county, from tributaries of the Rock, from the Du Page river near Joliet, from rocky spring branches in Union county and from springs in Calhoun and Jersey counties.

C. ricei (Nelson) *J JEC FR 327*

One specimen taken at a depth of 600 feet from Lake Michigan, in Grand Traverse bay off Old Mission, Michigan.

C. cognatus Richardson *J*

Cottus franklini Agassiz *JEC. Uranidea kumleinii* Hoy *FR 328*

EASTERN MILLER'S THUMB.—Three specimens taken in deep water from Lake Michigan, in Grand Traverse bay off Old Mission, Michigan.

GADIDAE

Lota (Cuvier) Oken

L. maculosa Le Sueur *J JEC FR 331*

LAWYER, BURBOT, LING.—Lake Michigan. Has also been taken from the Illinois at Peoria, Havana, Meredosia and

Naples, from the Rock at Milan and from the Mississippi at Rock Island.

NOTE

Melanotaenia Gill

M. nigrans Richardson

AUSTRALIAN RAINBOW FISH.—Three specimens were collected in July 1930 by Mr. Fred G. Orsinger of the John G. Shedd Aquarium, Chicago, Dr. Wilbur Luce of the University of Illinois and Dr. Thompson of the ILLINOIS STATE NATURAL HISTORY SURVEY. They were taken in a minnow seine, near the edge of a sand bar in the Mississippi river, out from the Southern Illinois Penitentiary. They were taken alive to the Shedd Aquarium, where they were identified by Dr. Carl Hubbs of the University of Michigan. It is supposed that they may have escaped from some tropical fish establishment in St. Louis, some 40 miles upstream.

BIBLIOGRAPHY

Forbes, Stephen Alfred and Robert Earl Richardson

- 1920 The fishes of Illinois. Natural History Survey of Illinois 3. cxxxvi+357 pp., 76 figs., 68 pls., 102 maps in separate atlas. \$4 plus postage on 6 lbs. ILLINOIS STATE NATURAL HISTORY SURVEY, Urbana.

Greene, C. Willard

- 1927 An ichthyological survey of Wisconsin. Papers of the Michigan Academy of Science, Arts and Letters 7(1926): 299.

Hubbs, Carl L.

- 1926 A check list of the fishes of the Great Lakes and tributary waters, with nomenclatorial notes and analytical keys. University of Michigan Museum of Zoology Miscellaneous Publications 15. 78 pp., 4 pls.

Hubbs, Carl L. and Laura C. Hubbs

- 1932 Experimental verification of natural hybridization between distinct genera of sunfishes. Papers of the Michigan Academy of Science, Arts and Letters 15(1931):427-37.

Jordan, David Starr

- 1929 Manual of the vertebrate animals of the northeastern United States inclusive of marine species. 446 pp. World Book Company, Yonkers-on-Hudson, New York.

Jordan, David Starr, Barton Warren Evermann and Howard Walton Clark

- 1930 Check list of the fishes and fishlike vertebrates of North and Middle America north of the northern boundary of Venezuela and Colombia. 670 pp. Report of the United States Commissioner of Fisheries for the Fiscal Year 1928, part 2. Document 1055.

Luce, Wilbur M.

- 1933 A survey of the fishery of the Kaskaskia river. ILLINOIS NATURAL HISTORY SURVEY BULLETIN 20(2):71-123. 12 figs. Fifty cents.

Thompson, David H.

- 1931 Variation in fishes as a function of distance. Transactions of the Illinois State Academy of Science 23(3):276-81. 2 figs.
- 1934 A concept of species among fishes. Transactions of the Illinois State Academy of Science 26(3):125.

Thompson, David H. and Francis D. Hunt

- 1930 The fishes of Champaign county. ILLINOIS NATURAL HISTORY SURVEY BULLETIN 19(1):5-101. 6 figs., 52 maps.

APPENDIX

HYBRIDIZATION AND RACIAL DIFFERENTIATION
AMONG ILLINOIS FISHES⁴

A list of the different kinds of Illinois fishes would be incomplete without mention of the hybrids which result from crosses between certain species.

Occurrence of Hybrids

Within certain groups of fishes and in certain localities, hybrids occur in frequencies as high as one in ten, or as low as one in several thousand. The greater number and variety of hybrids among these fishes, as compared with other groups of animals, may be a reflection of the breeding habit of these fishes, whereby the sperm of the male is set free in the water to be carried over wide areas while still alive and capable of fertilizing an egg. It seems somewhat more probable that in these fishes the devices for species recognition, or "consciousness of kind," are not so effective nor so rigidly fixed as among other animals. In this connection it may be mentioned that the groups of fishes which commonly produce natural hybrids are kinds (minnows, sunfishes and darters) which exhibit striking differences in the appearance of the two sexes. Those natural hybrids which have been studied in greatest detail are sterile, grow more rapidly than do the parent species, and show an excess of males. Two apparent exceptions to the rule of sterility will be mentioned later.

Experimental Confirmation

The bulk of our knowledge of natural hybrids among native fresh-water fishes has been contributed in recent years by Doctor Carl L. Hubbs. He has produced certain hybrid combinations of sunfishes in aquaria. Doctor Wilbur M. Luce during the last three seasons has developed a method for stripping and fertilizing eggs from sunfishes and has been able to rear the hybrid young in aquaria. Two of the

sunfishes listed as ordinary species in Forbes & Richardson (1920) have now been produced experimentally by crossing other native sunfishes. At the present time almost all of the hybrid combinations of the following six species of Illinois sunfishes have been found in Illinois waters: *Apomotis cyanellus*, *Allotis humilis*, *Xenotitis megalotis*, *Helioperca incisor*, *Eupomotis gibbosus* and *Chaenobryttus gulosus*.

The recognition of hybrids among these sunfishes is facilitated by the striking and distinctive color patterns of these species, certain characteristic features of both parents appearing in the hybrid as if the color patterns were superimposed. Hybrids between the pumpkinseed and the bluegill show two shapes of body as though they were reciprocal hybrids analogous to the mule and the hinny.

Other Common Hybrids

It is now reasonably certain that many minnows, previously regarded as anomalous and unclassifiable, are in reality species hybrids among various *Cyprinidae*. In Illinois, hybrids of the common shiner, *Luxilus cornutus*, with certain nest-building minnows (as pointed out by Doctor Hubbs), are especially common, presumably because of the fact that *Luxilus* habitually deposits its eggs in the nests of these other minnows. Hybrids within the group of darters are also believed to be rather common. A few readily recognized hybrids between goldfish and the European carp have been found in Illinois waters. Three or four catfishes have been taken in Rock river which appear to be hybrids between the channel cat, *Ictalurus punctatus*, and the flathead cat, *Opladelus olivaris*, both of which spawn in holes, hollow logs, etc. While it has not been proved that the so-called "mongrel" buffalo, *Ictiobus urus*, is really a hybrid

⁴By David H. Thompson.



Fig. 3.—Raising SURVEY hoop net along the east bank of the Illinois river below Havana.

between the other two native species, there is a considerable body of circumstantial evidence to indicate that such is the case. Some of the difficulties of identification in the genus *Carpiodes* and in the genus *Moxostoma* may be attributed to occasional hybridization within these genera.

Known Fertile Hybrids

In all of the above-named hybrids, whether proved or presumptive, the evidence indicates that they are in reality "mules" and incapable of propagation. In Illinois we have found two apparent exceptions to this rule. One of these has to do with the intergradation of *Cyprinella lutrensis* and *Cyprinella whipplii* in the region about the mouth of the Sangamon river near Chandlerville and Beardstown. At this point we not only find typical *lutrensis* and typical *whiplii* but also all possible combinations of the characteristics which are used to separate these two species.

Intergradations of this sort have also been found between the orange-spotted sunfish and the pumpkinseed sunfish in two localities—Senachwine lake near Henry and Crystal lake in Urbana. It seems certain that some hybrids between these two species have been fertile. Segregation and recombination of the characteristics of these two species have not been found in other waters of the state where they both occur. A rather detailed

study of the segregates of this pair of species in Lake Senachwine suggests that the hybrids between these two species have not bred *inter se* but have back-crossed to the orange-spotted sunfish.

Difference and Distance

A few years ago while making a detailed study of the fishes of the small streams which drain Champaign county, it was found that the fan-tailed darters, *Catnotus flabellaris*, of the streams draining into the Wabash had a consistently lower average number of spines in the dorsal fin than did those from streams draining into the Mississippi. It was also noticed that the orange-spotted sunfish, *Allotis humilis*, of the Sangamon river, which drains to the west, had a consistently steeper profile than did the same species in the Salt Fork, which drains to the east.

A study of the average dorsal spine and ray numbers of the Johnny darter, *Boleosoma nigrum*, collected from the various streams of Champaign county showed that the difference between the mean spine or ray numbers of any two populations increased with the water distance separating



Fig. 4.—Opening the tail compartment of a SURVEY hoop net, showing a catch of black crappies, carp and other fishes from the Illinois river.

these populations. The Johnny darter is well suited for such a study, since it is abundant, cosmopolitan in its range and sedentary in habit. The sedentary habit is to be emphasized because it appears that measurable differences between the mean spine or ray numbers of populations from different stream systems occur only among the more sedentary kinds. When such differences were sought among actively migrating kinds such as the suckers, they were not found. In brief, we may conclude that some kinds of fishes move so slowly that a measurable degree of evolution takes place while they are traveling from one stream to another.

Fin Phylogeny

A later and still more detailed study of the Johnny darters from all parts of Illinois, and using counts of the spines and rays of all of the fins, has substantiated the conclusion that the difference between the mean spine or ray numbers of two populations increases with increasing water distance separating these populations. This evidence indicates that these observed differences in average spine or ray numbers result, for the most part, from hereditary differences, and not from differences in the environment in which the various populations developed. The difference between the mean spine or ray numbers of any two populations is nearly proportional to the logarithm of water distance separating them.

It should be understood that water distance, as used here, implies water distance between different parts of the Mississippi system where there have been no barriers to the free movement of fishes until recent years. When the spine and ray numbers of the Johnny darter are compared with the spine and ray numbers of other subspecies of the Johnny darter, such as

Boleosoma nigrum olmstedii, (found in streams draining into the Atlantic from Quebec to Maryland) it is found that this difference is about twice as great as the average difference found between points within the Mississippi system separated by 3,000 miles of water.

Similarly, the amount of difference found between *Boleosoma nigrum* and another closely related Illinois species, *Vaillantia camura*, is about twice as great as the difference found between subspecies of the Johnny darter.

Other Taxonomic Differences

Not only are there consistent racial differences in the number of spines and rays of the Johnny darter and other sedentary fishes, but we also find consistent differences between different stream systems involving other taxonomic characteristics which are commonly used by ichthyologists to separate species, such as, for example, number of scales in the lateral line, number of scales bearing pores, number of rows of scales, color pattern, degree of scalation of body, cheek, nape and breast, the body contour and point of insertion of fins. Spine and ray numbers in the fins have been used in these studies merely because they seem to show nonadaptive variations, and because they may be easily and accurately treated in a quantitative manner. These considerations suggest that the degree of difference between any two fish populations forms a continuous series from the most insignificant racial differences up to the largest family differences.

Some of the evolutionary and genetic aspects of these racial differences in fishes as related to the nature of fish species have been discussed in two short papers published in the Transactions of the Illinois State Academy of Science (Thompson 1931, 1934).

INDEX TO FAMILIES, GENERA AND SPECIES

The index refers only to the mention or placement of names in the classified list, pp. 475-91. Synonymy is indicated by italic type. Family names are in capitals.

A

Abramis crysolencas 480
Acipenser fulvescens 475
rubicundus 475
 ACIPENSERIDAE 475
affinis, *Gambusia* 485
albus, *Parascaphirhynchus* 475
 alligator gar 477
Allotis humilis 487
Alosa ohioensis 477
alosoides, *Amphiodon* 477
Hiodon 477
alpenae, *Leucichthys* 478
Alvordius evermanni 488
maculatus 488
ouachitae 489
phoxocephalus 488
Ambloplites rupestris 487
amblops, *Hybopsis* 481
 AMBLYOPSIDAE 485
 AMEIURIDAE 483
Ameiurus lacustris 483
melas 484
natalis 484
nebulosus 484
 American eel 478
Amia calva 477
 AMIIDAE 477
Ammocrypta pellucida 488
Amphiodon alosoides 477
Anguilla bostoniensis 478
chrysypa 478
rostrata 478
anguilla, *Ictalurus* 483
 ANGUILLIDAE 478
anisurum, *Moxostoma* 479
annularis, *Pomoxis* 487
anogenus, *Hybopsis* 481
Notropis 481
anomalum, *Campostoma* 483
 APHREDODERIDAE 485
Aphredoderus sayanus 485
Aplodinotus grunniens 490
Apomotis cyanellus 486
euryorus 486
ischyrus 486
symmetricus 486
 appendix, *Lethenteron* 475
Argyrosomus artedi 478
artedi, *Argyrosomus* 478
Leucichthys 478
asprella, *Crystallaria* 488
aspro, *Hadropterus* 488
 ATHERINIDAE 486
atherinoides, *Notropis* 482
Atractosteus spatula 477
tristoechus 477
atripes, *Lythrurus* 482
Notropis umbratilis 482
atrocaudalis, *Hybopsis* 482
Notropis cayuga 482

atromaculatus, *Semotilus* 480
atronasus, *Rhinichthys* 480
auratus, *Carassius* 480
aureolum, *Moxostoma* 479
 Australian rainbow fish 491
 B
bairdii, *Cottus* 490
 banded darter 489
 barred darter 489
 bass, calico 487
 Kentucky 486
 large-mouth black 486
 rock 487
 small-mouth black 486
 southern small-mouth black 486
 warmouth 487
 white 490
 yellow 490
 big-eyed chub 481
 minnow 481
 big-mouth buffalo 478
 big-nosed sunfish 486
 big-toothed red horse 480
biguttatus, *Nocomis* 481
 billy gar 477
 black bass, large-mouth 486
 small-mouth 486
 southern small-mouth 486
 bullhead 484
 crappie 487
 horse 478
 lamprey, small 475
 sucker 479
 blackcheek top minnow 485
 blackfin 478, 482
 blackhead minnow 483
 black-nosed dace 480
 shiner 481
 black-sided darter 488
blennioides, *Diplesion* 488
Etheostoma 488
blennius, *Hybopsis* 482
Notropis 481
 blue cat 483
 herring 477
 sunfish 487
 blue-breasted darter 489
 bluefish 478
 bluegill 487
 blue-spotted sunfish 486
 blunt-nosed minnow 483
 river carp 479
Boleichthys fusiformis 489
Boleosoma camurum 488
nigrum 488
 boneless cat 475
 boops, *Hybopsis* 481
bostoniensis, *Anguilla* 478
 bowfin 477

bream 480
breviceps, *Moxostoma* 480
 brindled stonecat 484
 brook lamprey 475
 silverside 486
 stickleback 486
 trout, eastern 478
 brown bullhead 484
 trout, German 478
 brown-sided darter 489
bubalus, *Ictiobus* 478
buccata, *Ericymba* 483
 buffalo, big-mouth 478
 highback 479
 mongrel 478
 quillback 478
 razorback 479
 red-mouth 478
 round 478
 small-mouth 478
 bullhead, black 484
 brown 484
 minnow 483
 speckled 484
 yellow 484
 burbot 490

C

calico bass 487
calva, *Amia* 477
Campostoma anomalum 483
camura, *Vaillantia* 488
camurum, *Boleosoma* 488
Etheostoma 489
camurus, *Nothonotus* 489
canadense, *Cynoperca* 490
griseum, *Stizostedion* 490
canadensis grisea, *Cynoperca* 490
caprodes, *Percina* 488
Carassius auratus 480
carinatus, *Placopharynx* 480
 carp, blunt-nosed river 479
 common river 479
 European 480
 lake 479
 silver 479
 carpio, *Carpiodes* 479
Cyprinus 480
Carpiodes carpio 479
cyprinus 479
difformis 479
thompsoni 479
velifer 479
 cat, blue 483
 boneless 475
 channel 483
 eel 483
 flathead 484
 Fulton 483
 mud 484

silver 483
 spoonbill 475
 tadpole 484
 willow 483
 yellow 484
 cataractae, *Rhinichthys* 480
 catfish, northern 483
 of the lakes 483
Catonotus flabellaris 489
 CATOSTOMIDAE 478
Catostomus catostomus 479
 commersonii 479
 nigricans 479
 cave fish, spring 485
 Cayuga shiner 481
cayuga atrocaudalis, *Notropis* 482
 Hybopsis 481
 Notropis 481
 CENTRARCHIDAE 486
Centrarchus macropterus 488
 cepedianum, *Dorosoma* 477
Ceraticthys vigilax 483
Chaenobryttus gulosus 487
 channel cat 483
Chologaster papilliferus 485
Chrosomus erythrogaster 480
chrysochlorus, *Pomobolus* 477
Chrysoperca interrupta 490
 chrysops, *Lepibema* 490
 Rocca 490
chrysypa, *Anguilla* 478
 chub 478
 big-eyed 481
 creek 480
 flathead 481
 greaser 483
 river 481
 silver 481
 Storer's 481
 sucker 479
 cisco 478
 mooneye 478
Claricola squamiceps 489
Cliola vigilax 483
clupeaformis, *Coregonus* 477
 CLUPEIDAE 477
coeruleum, *Etheostoma* 489
coeruleus, *Oligocephalus* 489
cognatus, *Cottus* 490
commersonii, *Catostomus* 479
 common lake herring 478
 pike 485
 red horse 479
 river carp 479
 sculpin 490
 shiner 482
 sucker 479
 whitefish 477
 concolor, *Ichthyomyzon* 475
 COREGONIDAE 477
Coregonus clupeaformis 477
 quadrilateralis 477
 cornutus, *Luxilus* 482
 Notropis 482
 COTTIDAE 490
Cottogaster shumardi 488
Cottus bairdii 490

cognatus 490
franklini 490
ictalops 490
ricei 490
 crappie, black 487
 white 487
 creek chub 480
Cristivomer namaycush 478
crysoleucas, *Abramis* 480
 Notemigonus 480
Crystallaria asprella 488
cyanellus, *Apomotis* 486
 Lepomis 486
cycnocephalus, *Lythrurus um-*
 bratilis 482
Cycleptus elongatus 478
Cylindrosteus platostomus 477
Cynoperca canadense 490
 canadensis grisea 490
 grisea 490
Cyprinella lutrensis 482
 whippilii 482
cyprinella, *Ictiobus* 478
 Megastomatobus 478
 CYPRINIDAE 480
 CYPRINODONTIDAE 485
Cyprinus carpio 480
cyprinus, *Carpiodes* 479

D

dace, black-nosed 480
 horned 480
 long-nosed 480
 red-bellied 480
 darter, banded 489
 barred 489
 black-sided 488
 blue-breasted 489
 brown-sided 489
 fan-tailed 489
 fusiform 489
 green-sided 488
 Johnny 488
 least 490
 mud 489
 rainbow 489
 river 488
 sand 488
 sharp-nosed 488
 snub-nosed 488
deliciosus, *Hybopsis* 481
diaphanus menona, *Fundulus* 485
 Zygionectes 485
difformis, *Carpiodes* 479
Dionda nubila 483
Diplesion blennioides 488
dispar, *Fundulus* 485
 Zygionectes 485
dissimilis, *Erimystax* 481
 Hybopsis 481
 dogfish 477
dolomieu, *Micropterus* 486
Dorosoma cepedianum 477
 DOROSOMIDAE 477
dorsalis, *Hybopsis* 481

doughbelly 483
 drum, fresh-water 490
duquesnei, *Placopharynx* 480
 dwarf long-eared sunfish 487

E

eastern brook trout 478
 miller's thumb 490
 eel, American 478
 cat 483
 fresh-water 478
 lamper 475
Elassoma zonatum 488
 ELASSOMIDAE 488
elongatus, *Cycleptus* 478
emiliae, *Opsopoeodus* 480
Ericosma evides 489
Ericymba buccata 483
Erimystax dissimilis 481
Erimyzon oblongus 479
 sucetta 479
 oblongus 479
Erinemus hyalinus 481
 storerianus 481
Erogala whippilii 482
erythrogaster, *Chrosomus* 480
 ESOCIDAE 484
Esox estor 485
 lucius 485
 masquinongy 485
 vermiculatus 484
estor, *Esox* 485
Etheostoma blennioides 488
 camurum 489
 coeruleum 489
 flabellare 489
 iowae 489
 jessiae 489
 obeyense 489
 squamiceps 489
 zonale 489
 ETHEOSTOMIDAE 488
Eucalia inconstans 486
Eupomotis gibbosus 487
 heros 487
 European carp 480
euryorus, *Apomotis* 486
 Lepomis 486
evermanni, *Alvordius* 488
 Hadropterus 488
evides, *Ericosma* 489
 Hadropterus 489
exilis, *Oligocephalus* 489
 Rabida 484
 Schilbeodes 484
Extrarius hyostomus 480

F

fan-tailed darter 489
 fathead 483
 fiddler 483
 fine-scaled sucker 479
 fish, Australian rainbow 491
 mosquito 485
 spring cave 485
flabellare, *Etheostoma* 489

flabellaris, *Catnotus* 489
 flathead cat 484
 chub 481
 flavescens, *Perca* 490
 flavus, *Noturus* 484
 flier 488
floridana, *Huro* 486
 fontinalis, *Salvelinus* 478
 Forbesella papillifera 485
franklini, *Cottus* 490
 freckled stonecat 484
 fresh-water drum 490
 eel 478
 Fulton cat 483
 fulvescens, *Acipenser* 475
 Fundulus diaphanus menona
 485
 dispar 485
 notatus 485
 furcatus, *Ictalurus* 483
 fusiform darter 489
 fusiformis, *Boleichthys* 489
 Hololepis 489

G

GADIDAE 490
gairdnerii, *Salmo* 478
Gambusia affinis 485
 patruelis 485
 gar, alligator 477
 billy 477
 long-nosed 477
 short-nosed 477
 GASTEROSTEIDAE 486
gelida, *Macrhybopsis* 480
gelidus, *Macrhybopsis* 480
 German brown trout 478
gibbosus, *Eupomotis* 487
gilberti, *Hybopsis* 481
 Notropis 481
 gizzard shad 477
 goggle-eye 487
 golden shad 477
 shiner 480
 goldeye 477
 goldfish 480
 goujon 484
gracilis, *Platygobio* 481
 grass pike 484
 gray pike 490
 greaser chub 483
 great northern pike 485
 green sunfish 486
 green-sided darter 488
 grindle 477
 grisea, *Cynoperca* 490
 canadensis 490
 Cynoperca 490
griseum, *Stizostedion canadense*
 490
 grunniens, *Aplodinotus* 490
 gulosus, *Chaenobryttus* 487
guttatus, *Percopsis* 485
 gyrinus, *Schilbeodes* 484

H

hackleback 475

Hadropterus aspro 488
 evermanni 488
 evides 489
 ouachitae 489
 phoxocephalus 488
 scierus 489
 Hagochila lacera 480
 hammerhead 479
Haustor lacustris 483
Helioperca incisor 487
 heros, *Eupomotis* 487
 herring, blue 477
 common lake 478
 lake 478
 toothed 477
 heterodon, *Hybopsis* 481
 Notropis 481
 heterolepis, *Hybopsis* 481
 hickory shad 477
 highback buffalo 479
Hiodon alosoides 477
 tergisus 477
 HIODONTIDAE 477
 hog sucker 479
Hololepis fusiformis 489
 horned dace 480
 hornyhead 481
 horse, big-toothed red 480
 black 478
 common red 479
 short-headed red 480
 hoyi, *Leucichthys* 478
Hudsonius hudsonius 482
hudsonius, *Hybopsis* 482
 Notropis 482
 humilis, *Allotis* 487
 Lepomis 487
Huro floridana 486
 salmoides 486
 hyalinus, *Erinemus* 481
Hybognathus nubila 483
 nuchalis 483
Hybopsis amblops 481
 anogenus 481
 atrocaudalis 482
 blennius 482
 boops 481
 cayuga 481
 deliciosus 481
 dissimilis 481
 dorsalis 481
 gilberti 481
 heterodon 481
 heterolepis 481
 hudsonius 482
 hyostomus 480
 kentuckiensis 481
 nux richardsoni 482
 phenacobius 481
 shumardi 481
 storerianus 481
 volucellus 481
Hyborhynchus notatus 483
hyostoma, *Macrhybopsis* 480
hyostomus, *Extrarius* 480
 Ilybopsis 480
Hypentelium nigricans 479

I

Ichthyomyzon concolor 475
ictalops, *Cottus* 490
Ictalurus anguilla 483
 furcatus 483
 punctatus 483
Ictiobus bubalus 478
 cyprinella 478
 urus 478
illecebrosus, *Notropis* 481
Imostoma shumardi 488
incisor, *Helioperca* 487
inconstans, *Eucalia* 486
interrupta, *Chrysoperca* 490
 Morone 490
iowae, *Etheostoma* 489
 Oligocephalus 489
iridea, *Trutta* 478
irideus, *Salmo* 478
ischyrus, *Apomotis* 486
 Lepomis 486

J

jack salmon 490
jejunus, *Notropis* 482
 Paranotropis 482
jessiae, *Etheostoma* 489
 Oligocephalus 489
johannae, *Leucichthys* 478
 Johnny darter 488

K

kentuckiensis, *Hybopsis* 481
 Nocomis 481
 Kentucky bass 486
 killifish, *Menona* 485
kiyi, *Leucichthys* 478
kumleinii, *Uranidea* 490

L

Labidesthes sicculus 486
lacera, *Hagochila* 480
lacustris, *Ameiurus* 483
 Haustor 483
 Villarius 483
 lake carp 479
 herring 478
 shiner 482
 sturgeon 475
 trout 478
 lakes, catfish of the 483
 lamper 475
 eel 475
Lampetra wilderi 475
 lamprey 475
 brook 475
 small black 475
 landlocked salmon 478
 large-mouth black bass 486
 lawyer 490
 least darter 490
Lepibema chrysops 490
 LEPISOSTEIDAE 477
Lepisosteus osseus 477

platostomus 477
tristoechus 477
Lepomis cyanellus 486
euryor 486
humilis 487
ischyrus 486
megalotis 487
peltastes 487
miniatus 487
pallidus 487
symmetricus 486
Leptops olivaris 484
lesueurii, *Moxostoma* 480
Lethenteron appendix 475
Lethogrammus symmetricus 486
Leucichthys alpenae 478
artedi 478
hoi 478
johannae 478
kiyi 478
nigripinnis 478
reighardi 478
zenithicus 478
limi, *Umbra* 484
ling 490
little pickerel 484
log perch 488
long-eared sunfish 487
dwarf 487
longjaw 478
long-nosed dace 480
gar 477
sturgeon 475
sucker 479
Lota maculosa 490
lucius, *Esox* 485
lutrensis, *Cyprinella* 482
Moniana 482
Notropis 482
Luxilus cornutus 482
Lythrurus atripes 482
umbratilis 482
cynocephalus 482

M

Macrhybopsis gelida 480
gelidus 480
hyostoma 480
macropterus, *Centrarchus* 488
maculatus, *Alvordius* 488
maculosa, *Lota* 490
mad Tom 484
masquinongy, *Esox* 485
megalotis, *Lepomis* 487
peltastes, *Lepomis* 487
Xenotis 487
Megastomatobus cyprinella 478
melanops, *Minytrema* 479
Melanotaenia nigrans 491
melas, *Ameiurus* 484
Menominee whitefish 477
Menona killifish 485
top minnow 485
menona, *Fundulus diaphanus* 485

Zygonectes diaphanus 485
Microperca punctulata 490
micropogon, *Nocomis* 481
Micropterus dolomieu 486
pseudaplites 486
salmoides 486
miller's thumb 490
eastern 490
miniatus *Lepomis* 487
Sclerotis 486
minnow, *big-eyed* 481
blackcheek top 485
blackhead 483
blunt-nosed 483
bullhead 483
Menona top 485
mud 484
silver-mouth 483
silvery 483
spot-tailed 482
steel-colored 482
straw-colored 481
sucker-mouth 483
top 485
viviparous top 485
Minytrema melanops 479
mirabilis, *Phenacobius* 483
Missouri sucker 478
miurus, *Rabida* 484
Schilbeodes 484
mongrel buffalo 478
Miniana lutrensis 482
mooneye 477, 478
cisco 478
northern 477
Morone interrupta 490
MORONIDAE 490
mosquito fish 485
Moxostoma anisurum 479
aureolum 479
breviceps 480
lesueurii 480
mud cat 484
darter 489
minnow 484
mudfish 484
muskellunge 485

N

namaycush, *Cristivomer* 478
Nanostoma zonale 489
natalis, *Ameiurus* 484
nebulosus, *Ameiurus* 484
nigrans, *Melanotaenia* 491
nigrans, *Catostomus* 479
Hypentelium 479
nigripinnis, *Leucichthys* 478
nigrum, *Boleosoma* 488
nine-spine stickleback 486
Nivicola obeyense 489
Nocomis biguttatus 481
kentuckiensis 481
micropogon 481
nocturna, *Rabida* 484
nocturnus, *Schilbeodes* 484
northern catfish 483
mooneye 477
pike, *great* 485

sucker 479
notatus, *Fundulus* 485
Hyborhynchus 483
Pimephales 483
Zygonectes 485
Notemigonus crysoleucas 480
Nothonotus camurus 489
Notropis anogenus 481
atherinoides 482
blennius 481
cayuga 481
atrocaudalis 482
cornutus 482
gilberti 481
heterodon 481
hudsonius 482
illecebrosus 481
jejunus 482
lutrensis 482
phenacobius 481
photogenis 482
pilsbryi 482
rubrifrons 482
umbratilis atripes 482
whippilii 482
Noturus flavus 484
nubila, *Dionda* 483
Hybognathus 483
nuchalis, *Hybognathus* 483
nux richardsoni, *Hybopsis* 482

O

obeyense, *Etheostoma* 489
Nivicola 489
obeyensis, *Oligocephalus* 489
oblongus, *Erimyzon* 479
Erimyzon sucetta 479
ohiensis, *Alosa* 477
Ohio shad 477
old-fashioned sunfish 487
Oligocephalus coeruleus 489
exilis 489
iowae 489
jessiae 489
obeyensis 489
olivaris, *Leptops* 484
Opladelus 484
omiscomaycus, *Percopsis* 485
Opladelus olivaris 484
Opsopoeodus emiliae 480
orange-spotted sunfish 487
osseus, *Lepisosteus* 477
ouachitae, *Alvordius* 489
Hadropterus 489

P

paddlefish 475
pallidus, *Lepomis* 487
papillifera, *Forbesella* 485
papilliferus, *Chologaster* 485
Paranotropsis jejunos 482
Parascaphirhynchus albus 475
patruelis, *Gambusia* 485
pellucida, *Ammocrypta* 488
pellucidus, *Vigil* 488
peltastes, *Lepomis megalotis* 487

Xenotis megalotis 487

Perca flavescens 490

perch, log 488

pirate 485

red 487

ringed 490

trout 485

white 490

yellow 490

PERCIDAE 490

Percina caprodes 488

PERCOPSIDAE 485

Percopsis guttatus 485

omiscomaycus 485

PETROMYZONIDAE 475

Phenacobius mirabilis 483

phenacobius, *Hybopsis* 481

Notropis 481

photogenis, *Notropis* 482

phoxocephalus, *Alvordius* 488

Hadropterus 488

pickerel 485

little 484

pigmy sunfish 488

pike, common 485

grass 484

gray 490

great northern 485

sand 490

wall-eyed 490

pilsbryi, *Notropis* 482

Pimephales notatus 483

promelas 483

pirate perch 485

Placopharynx carinatus 480

duquesnei 480

platyrhynchus, *Scaphirhynchus* 475

platostomus, *Cylindrosteus* 477

Lepisosteus 477

Platygobio gracilis 481

POECILIIDAE 485

Polyodon spathula 475

POLYODONTIDAE 475

Pomobolus chrysochlorus 477

Pomoxis annularis 487

sparoides 487

promelas, *Pimephales* 483

Prosopium quadrilaterale 477

pseudaplites, *Micropterus* 486

pugnose shiner 481

pumpkinseed 487

punctatus, *Ictalurus* 483

punctulata, *Microperca* 490

Pungitius pungitius 486

Pygosteus pungitius 486

Q

quadrilaterale, *Prosopium* 477

quadrilateralis, *Coregonus* 477

quillback 479

buffalo 478

R

rabbit-mouth sucker 480

Rabida exilis 484

miurus 484

nocturna 484

rainbow darter 489

fish, Australian 491

trout 478

razorback buffalo 479

red horse, big-toothed 480

common 479

short-headed 480

perch 487

sucker 479

red-bellied dace 480

redfin 482

red-mouth buffalo 478

reighardi, *Leucichthys* 478

Rhinichthys atronasmus 480

cataractae 480

ricei, *Cottus* 490

richardsoni, *Hybopsis nux* 482

ringed perch 490

river carp, blunt-nosed 479

common 479

chub 481

darter 488

roach 480

Roccus chrysops 490

rock bass 487

sturgeon 475

rostrata, *Anguilla* 478

rosy-faced shiner 482

round buffalo 478

sunfish 488

whitefish 477

rubicundus, *Acipenser* 475

rubrifrons, *Notropis* 482

rupestris, *Ambloplites* 487

S

salar sebago, *Salmo* 478

Salmo gairdnerii 478

irideus 478

salar sebago 478

salmoides, *Huro* 486

Micropterus 486

salmon, jack 490

landlocked 478

Sebago 478

SALMONIDAE 478

Salvelinus fontinalis 478

sand darter 488

pike 490

sauger 490

sayanus, *Aphredoderus* 485

Scaphirhynchus platyrhynchus

475

Schilbeodes exilis 484

gyrinus 484

miurus 484

nocturnus 484

SCIAENIDAE 490

sciera, *Serraria* 489

scierus, *Hadropterus* 489

Sclerotis miniatus 486

sculpin, common 490

Sebago salmon 478

sebago, *Salmo salar* 478

Semotilus atromaculatus 480

Serraria sciera 489

shad 479

gizzard 477

golden 477

hickory 477

Ohio 477

sharp-nosed darter 488

sheepshead 490

shiner, black-nosed 481

Cayuga 481

common 482

golden 480

lake 482

pugnose 481

rosy-faced 482

silver 482

spotted 481

short-headed red horse 480

short-nosed gar 477

shovel-nosed sturgeon 475

shumardi, *Cottogaster* 488

Hybopsis 481

Imostoma 488

sicculus, *Labidesthes* 486

silver carp 479

cat 483

chub 481

shiner 482

silvereen 479

silver-mouth minnow 483

silverside, brook 486

silvery minnow 483

skipjack 477

slender stonecat 484

small black lamprey 475

small-mouth black bass 486

southern 486

buffalo 478

snub-nosed darter 488

southern small-mouth black

bass 486

sparoides, *Pomoxis* 487

spathula, *Polyodon* 475

spatula, *Atractosteus* 477

speckled bullhead 484

trout 478

spoonbill cat 475

spot-tailed minnow 482

spotted shiner 481

sucker 479

spring cave fish 485

squamiceps, *Claricola* 489

Etheostoma 489

steel-colored minnow 482

stickleback, brook 486

nine-spine 486

Stizostedion canadense griseum

490

vitreum 490

stonecat 484

brindled 484

freckled 484

slender 484

stoneroller 479, 483

storerianus, *Erinemus* 481

Hybopsis 481

Storer's chub 481

straw-colored minnow 481

striped sucker 479

sturgeon, lake 475
 long-nosed 475
 rock 475
 shovel-nosed 475
 white 475
 sucetta, *Erimyzon* 479
 oblongus, *Erimyzon* 479
 sucker, black 479
 chub 479
 common 479
 fine-scaled 479
 hog 479
 long-nosed 479
 Missouri 478
 northern 479
 rabbit-mouth 480
 red 479
 spotted 479
 striped 479
 sweet 479
 white-nosed 479
 sucker-mouth minnow 483
 sunfish, big-nosed 486
 blue 487
 blue-spotted 486
 dwarf long-eared 487
 green 486
 long-eared 487
 old-fashioned 487
 orange-spotted 487
 pigmy 488
 round 488
 Superior whitefish 477
 sweet sucker 479
 symmetricus, *Apomotis* 486
 Lepomis 486
 Lethogrammus 486

T

tadpole cat 484
 tergisus, *Hiodon* 477
 thompsoni, *Carpiodes* 479

thumb, miller's 490
 eastern miller's 490
 Tom, mad 484
 toothed herring 477
 top minnow 485
 blackcheek 485
 Menona 485
 viviparous 485
 topwater 485
 tristoecus, *Atractosteus* 477
 Lepisosteus 477
 trout, eastern brook 478
 German brown 478
 lake 478
 perch 485
 rainbow 478
 speckled 478
 Trutta iridea 478
 trutta 478

U

Umbra limi 484
 umbratilis *atripes*, *Notropis* 482
 Cyanocephalus, *Lythrurus* 482
 Lythrurus 482
 UMBRIDAE 484
Uranidea kumleini 490
 urus, *Ictiobus* 478

V

Vaillantia camura 488
velifer, *Carpiodes* 479
 vermiculatus, *Esox* 484
 Vigil pellucidus 488
 Vigilax, *Ceraticthys* 483
 Cliola 483
 Villarius lacustris 483
 vitreum, *Stizostedion* 490
 viviparous top minnow 485
 volucellus, *Hybopsis* 481

W

walleye 490
 wall-eyed pike 490
 warmouth bass 487
 whippilii, *Cyprinella* 482
 Erogala 482
 Notropis 482
 white bass 490
 crappie 487
 perch 490
 sturgeon 475
 whitefish, common 477
 Menominee 477
 round 477
 Superior 477
 white-nosed sucker 479
wilderi, *Lampetra* 475
 willow cat 483

X

Xenotis megalotis 487
 peltastes 487

Y

yellow bass 490
 bullhead 484
 cat 484
 perch 490

Z

zenithicus, *Leucichthys* 478
 zonale, *Etheostoma* 489
 Nanostoma 489
 zonatum, *Elassoma* 488
Zygonectes diaphanus menona 485
 dispar 485
 notatus 485

INDEX

The reader is referred to preceding indexes of families, genera and species at pages 270-79 for Dermaptera and Orthoptera, 468-71 for Plecoptera, and 495-500 for the fishes of Illinois. Inasmuch as articles I and II were printed without indexes, items pertaining to them have been gathered below under the headings "elm disease" and "fishery of the Kaskaskia river," respectively.

A

Ambraw river—see Embarrass river
Anax, facing 473, 474
Apple river 476
Australian rainbow fish 491

B

bacteria on elm 58
Big Bay creek 476
Big Muddy river 476

C

Cache river 476
canal, Illinois-Mississippi 476
chief plant inspector, requests elm disease investigation 3
cockroaches—see Orthoptera
control 269
Crooked creek 476

D

Dermaptera 150, 151
Desplaines river 476
Diseases of elm—see elm disease

E

elm disease 1-70
agars 13, 15, 32-34, 38, 40-45, 48, 50-57
Alternaria 17, 19, 20, 44, 45, 51, 52
American elm, anthracnose in 1
investigation of 1
leaf spot 1
Anthostomella 17, 19, 41, 49, 52
anthracnose in American elm 1
ascomycetous fungus 52
bacteria 58
Basisporium gallarum 54
Buisman, examines Illinois trees 1
cankers 8, 10, 11, 16, 18, 25, 41-43, 45, 46, 52
cause, thought to be elm borer 3
Cephalosporium 17, 19, 51
Ceratostomella Ulmi 2
Champaign, Illinois, *Vermicularia* in 39
chief plant inspector, requests investigation 3
combinations of fungi in isolations 19

elm disease—cont'd

Coniothyrium 17-32, 37-40, 42, 43, 49, 59, 65-67

Fuckelii 37

Ulmi 37

contamination, culture 17

control experiments 59-67

cultures 12

contamination 17

general results of tests 15-20

stock 13

Cytospora 17, 20, 52, 53

ambicus 53

Danville, Illinois, *Coniothyrium* in 21
data, recording 14

Diaporthe 46

Diplodia 17, 20, 53, 54

Ulmi 53, 54

distribution, statewide 9

Dutch elm disease, caused by *Ceratostomella Ulmi* 2

found by May in Ohio 1

not in Illinois 2

presence suspected 1

dying elms, observed prior to 1883 3

El Dara, Illinois, *Coniothyrium* in 21

elm borer, thought cause of disease 3

"elm wilt," erroneous name 1, 20

elms examined, wild 11

Endothia 13

Epicoccum 17, 19, 20, 57, 58

experiments, control 59-67

Forbes, Dr. S. A., observes dying elms
prior to 1883 3

Freeman nursery, cooperates in study
2, 61

fungous parasitization, symptom of 6
fungi, ascomycetous 52

combinations of in isolations 19

culturing 12, 13

genera recognized 17

isolations 12, 17

nonfruiting 17

parasitic 19, 20, 50, 59

elm disease—cont'd

- fungi, saprophytic 19, 20, 45, 55, 59
 - species recognized 17
 - 200 specimens examined 15
- Fusarium* 2, 17, 20, 41, 50-52
 - oxysporum* 50
 - scirpi compactum* 50
- genera of fungi isolated 17
- general symptoms 6
- Gnomonia ulmea* 1
- Graphium Ulmi* 2, 58
- gunmosis 28
- Helminthosporium* 17, 57, 58
- histological studies 13
- Illinois, previous outbreaks in 3
- Illinois State Nurserymen's Association, cooperates in study 2
- importance of disease, in nurseries 9
 - in cities 9
- infections, other than specific 50-58
- inoculation trials 58, 59
- investigation, methods of 12
 - of American elm 1
 - requested by chief plant inspector 3
- isolations, of fungi 12, 17
 - combinations of fungi in 19
 - 512 made 15
 - percentage sterile 18
 - 20 genera recognized in 15
- La Grange, Illinois, *Anthostomella*, *Fusarium*, *Nigrospora*, *Vermicularia* in 40, 41
- late-season symptoms 9
- leaf spot of American elm 1
- Libertyville, Illinois, *Coniothyrium* in 21
- manifestations of disease 20
- May, finds Dutch elm disease in Ohio 1
- methods of investigation 12
- Micrococcus Ulmi* 58
- Mounds City, Illinois, *Coniothyrium* in 21
- Mycosphaerella Ulmi* 1
- Nigrospora*, 17, 19, 41, 54, 55
 - sphaerica* 54, 55
- nonfruiting fungi 17
- nurseries, Freeman 61
 - Swain-Nelson 61
- nursery stock, symptoms in 6
- nurserymen's association, cooperates 2

elm disease—cont'd

- Ohio, Dutch elm disease in 1
- other infections 50-58
- outbreak, in Illinois 3
 - in Kentucky 4
 - in Massachusetts 4
- parasitic fungi 19, 20, 50, 59
- Phoma* 17-20, 41-45, 49, 59, 67
 - alternariaceum* 45
- Phomopsis* 17, 20, 45, 46, 48
- Pianeze III-b stain 13
- previous outbreaks 3, 4
- pruning 61-66
- Pseudomonas lignicola* 58
- raspberry, *Coniothyrium Fuckelii* on 37, 38
- recording data 14
- results of cultural tests 15
- Rhabdospora* 17, 57, 58
- Rockford, Illinois, unthrifty elms in 4
- Saperda tridentata*, thought to be cause of disease 3
- saprophytes 19, 20
- source of specimens 11, 15
- Sphaeropsis* 17, 18, 20, 45, 67
 - ulmicola* 45
- species of fungi recognized 17
- specific diseases 20-50
 - symptoms 8
- specimens, source of 11, 15
- spraying 61-66
- stain, Pianeze III-b 13
- state entomologist, observes dying elms prior to 1883 3
- statewide distribution of disease 9
- Stemphylium* 17-20, 51, 52
- sterile isolations 18
- stock cultures 13
- Swain-Nelson nursery, cooperates in study 2, 61
- symptoms, general 6
 - in nursery stock 6
 - of fungous parasitization 6
 - specific 8
 - late-season 9
- tannins 33
- Thielavia* 13
- tracheomycosis, associated with *Coniothyrium* spp. 20-38, 39
- treatment, heretofore unable to recommend 9

elm disease—concl'd

Ulmus americana 1, 37*campestris* 37unthrifty elms, in Rockford, Illinois +
Urbana, Illinois, *Coniothyrium*, *Vermicularia* in 39*Vermicularia* 17-19, 35, 36, 38-41, 67*Vorticillium* 17-19, 47-50, 67*alboatrum* 48Westerdijk and Buisman, 1929 in-
vestigations 1

wild elms examined 11

Xylaria 17, 55-57

Embarrass river 476

F

fishery of the Kaskaskia river 71-123

Abramis crysolencas 115abundance of fishes, factors affecting
89

age determination of fishes 94

air and water temperatures 75

alligator gar 97

Allotis humilis 110*Alzordius maculatus* 119*phoxocephalus* 119*Ameirus melas* 107*natalis* 107*nebulosus* 107*Amia calva* 98*Ammocrypta pellucida* 120*Anguilla hostoniensis* 99*chrysypa* 99*Aphredoderus sayanus* 121*Aplodinotus grunniens* 113*Atractosteus tristoechus* 97

bass, large-mouth black 109

rock 94, 112

small-mouth black 109

warmouth 111

big-eyed chub 117

minnow 116

black bass, large-mouth 109

small-mouth 109

bullhead 72

crappie 113

sucker 101

blackfin minnow 115

black-head minnow 118

black-sided darter 119

blue cat 96, 105, 109

fishery of the Kaskaskia river—cont'd

bluegill 111

blunt-nosed minnow 118

Boleichthys fusiformis 120*Bolcosoma camurum* 120*nigrum* 120

boneless cat 96

bottom fauna 89

bowfin 98

brindled stonecat 119

brook lamprey 95

silverside 121

buffalo 73, 84, 88, 95, 100, 101

bullhead 73, 87, 107

black 72

minnow 117

Campostoma anamolium 118

carp 73, 84, 88, 95, 103

suckers 101

Carpiodes carpio 101*cyprinus* 101*difformis* 101*velifer* 101

cat, blue 96, 105, 109

boneless 96

channel 73, 94, 96, 106, 109

flathead 73

Fulton 73

mud 94, 108

spoonbill 73, 95, 96

tadpole 118

catches 87, 88

value 83-85

catfish 72, 84, 87, 88, 95, 118

migrations 72

Catnotus flabellaris 120*Catostomus commersonii* 101*nigricans* 101

caviar, spoonbill roe 95

Ceraticthys vigilax 117*Chaenobryttus gulosus* 111

channel cat 73, 94, 96, 106, 109

chub, big-eyed 117

flat-headed 116

Kentucky 116

Storer's 116

sucker 102

Cliala vigilax 117

collection of fishes by S. A. Forbes 71

commercial fishing 71, 73, 84, 91

common red horse 102

fishery of the Kaskaskia river—cont'd

common shiner 115
 sucker 101
Cottogaster shumardi 119
 crappie, black 113
 white 74, 112
 creeks 79
Cylindrostens platostomus 97
Cynoperca canadensis 109
Cyprinella lutrensis 116
 whippelii 116
Cyprinus carpio 103
 dace, horned 115
 darter, black-sided 119
 fantail 120
 Johnny 120
 rainbow 120
 sand 120
 sharp-nosed 119
 snub-nosed 120
 direction of flow 74
 discharge 78
 disease of fishes 91
 dogfish 98
Dorosoma cepedianum 99
 doughbelly 118
 drainage basin 77, 92, 93
 drouth, recovery from 72
 drum 84, 113
 eel 99
 elevation 74
 enemies of fishes 91
 equipment and methods 80
Ericosma evides 119
Ericymba buccata 117
Erimystax dissimilis 117
Erimyzon succetta oblongus 102
Erinemus hyalinus 117
 storerianus 116
Esox vermiculatus 108
Etheostoma coeruleum 120
 flabellare 120
 jessiae 120
Extrarius hyostomus 117
 factors affecting abundance of fishes
 89
 fantail darter 120
 fathead minnow 118
 fauna, bottom 89
 fishes, age determination of 94
 disease of 91

fishery of the Kaskaskia river—cont'd

fishes, enemies of 91
 factors affecting abundance of 89
 hybrid 115, 116
 kinds of caught 87
 large 95-114
 of Mississippi river in Kaskaskia 73
 rate of growth 93
 small 114-21
 tagged 72
 fishing, clubs along Kaskaskia 84
 commercial 71, 73, 84, 91
 hook and line 71, 73, 91
 flathead cat 73
 flat-headed chub 116
 floodplain 79
 flow, direction of 74
 Forbes, S. A., collection of fishes by 71
 freckled stonecat 118
 frogs and turtles 83
 Fulton cat 73
Fundulus notatus 120
Gambusia affinis 121
 patruclis 121
 gar, alligator 97
 long-nosed 97
 short-nosed 97
 gizzard shad 99
 glaciation 76, 77
 golden shiner 115
 goujon 108
 gradient 75
 grass pike 108
 grindle 98
 grinnel 98
 growth rate of fishes 93
Hadropterus aspro 119
 evides 119
 phoxocephalus 119
 headquarters 80
Helioperca incisor 111
 herring, toothed 99
Hexagenia, food of spoonbill 95
Hiodon tergisus 99
 hog sucker 101
 hook and line fishing 71, 73, 91
 hoop-net catches 87, 88
Hololepis fusiformis 120
 horned dace 115
 hornyhead 116
Huro salmoides 109

fishery of the Kaskaskia river—cont'd

Hybognathus nuchalis 117
Hybopsis amblops 117
blennius 116
boops 116
deliciosus 116
dissimilis 117
gelidus 117
hudsonius 116
hyostomus 117
kentuckiensis 116
storerianus 116
Hyborhynchus notatus 118
 hybrids 115, 116
 hydrography 74
Hypentelium nigricans 101
Ichthyomyzon concolor 95
Ictalurus furcatus 105
punctatus 106
Ictiobus bubalus 100
cyprinella 100
urus 100
 Illinois, glaciation 76, 77
 main rivers of 83
Imostoma shumardi 119
 information, sources of 83
 investigation, methods and equipment 80
 personnel 80
 Johnny darter 120
 Kaskaskia, Illinois 73, 80
 river 74, 77
 Kentucky chub 116
 kinds of fishes caught 87
 of tackle 88
Labidesthes sicculus 121
 lake lamprey 95
 lamprey, brook 95
 lake 95
 silvery 95
 large fishes 95-114
 large-mouth black bass 109
 length of Kaskaskia river 74
Lepisosteus osseus 97
platostomus 97
tristoechus 97
Lepomis humilis 110
megalotis 111
pallidus 111
Leptops olivaris 108
Lethenteron appendix 95

fishery of the Kaskaskia river—cont'd

long-eared sunfish 111
 long-nosed gar 97
 log perch 119
Luxilus cornutus 115
 × *Hybognathus nuchalis* 116
 × *Nocomis biguttatus* 115
Lythrurus umbratilis cyanocephalus 115
Macrhybopsis gelidus 117
 mad Tom 118
 main rivers of Illinois 83
Megastomatobus cyprinella 100
 methods and equipment 80
Micropterus dolomieu 109
salmoides 109
 migrations 72
 minnow, big-eyed 116
 blackfin 115
 black-head 118
 blunt-nosed 118
 bullhead 117
 fathead 118
 silver-mouthed 117
 silvery 117
 snub-nosed 115
 spot-tailed 116
 straw-colored 116
 sucker-mouthed 117
 top 120
 viviparous top 121
Minytrema melanops 102
 miscellaneous small fishes 120
 Mississippi river 79
 fishes in Kaskaskia 73
 moon eye 99
 moraine, Shelbyville 75
 mouth of Kaskaskia river 74
Moxostoma aureolum 102
breviceps 103
 mud cat 94, 108
 mussels, pearls and slugs 83-86
Nocomis biguttatus 116
Notemigonus crysoleucas 115
Notropis antherinoides 115
 cornutus 115
 blennius 116
 hudsonius 116
 illecebrosus 116
 jejunus 116
 lutrensis 116

fishery of the Kaskaskia river—cont'd

- Notropis pilsbryi* 115
- umbratilis atripes* 115
- whifflii* 116
- Noturus flavus* 118
- Okaw river 71
- Oligocephalus coeruleus* 120
- jessiae* 120
- Opladelus olivaris* 108
- Opsopocodus emiliae* 115
- orange-spotted sunfish 110
- paddlefish 96
- pearls, slugs and mussels 83-86
- perch, log 119
- pirate 121
- Percina caprodes* 119
- personnel of investigation 80
- Petromyzon marinus dorsatus* 95
- Phenacobius mirabilis* 117
- pike, grass 108
- wall-eyed 109
- Pimephales notatus* 118
- promelas* 118
- pirate perch 121
- plankton 90
- sampling 82
- Platygobio gracilis* 116
- pollution 92
- Polyodon spathula* 96
- Pomoxis annularis* 112
- sparoides* 113
- Rabida exilis* 119
- miurus* 119
- rainbow darter 120
- rate of growth in fishes 93
- recovery from drouth 72
- red horse, common 102
- short-headed 103
- redfin 116
- river, Mississippi 73, 79
- Okaw 71
- Sangamon 74
- shiner 115
- rivers of Illinois, main 83
- rock bass 94, 112
- sand darter 120
- Sangamon river, white crappie in 74
- Sargasso sea, spawning ground of eel 100
- sauger 109

fishery of the Kaskaskia river—cont'd

- Schilbeodes exilis* 119
- gyrinus* 118
- miurus* 119
- nocturnus* 118
- Semotilus atromaculatus* 115
- shad, gizzard 99
- sharp-nosed darter 119
- sheepshead 84, 113
- Shelbyville moraine 75
- shiner, common 115
- golden 115
- river 115
- spotted 117
- short-headed red horse 103
- short-nosed gar 97
- silverfin 116
- silver-mouthed minnow 117
- silverside, brook 121
- silvery lamprey 95
- minnow 117
- slender stonecat 119
- slugs, mussels and pearls 83-86
- small fishes 114-21
- small-mouth black bass 109
- smoked sturgeon 96
- snub-nosed darter 120
- minnow 115
- source of Kaskaskia 74
- sources of information on fishery 83
- spoonbill cat 73, 95, 96
- spot-tailed minnow 116
- spotted shiner 117
- sucker 102
- Stizostedion canadense griseum* 109
- vitreum* 109
- stonecat, brindled 119
- freckled 118
- slender 119
- stoneroller 118
- Storer's chub 116
- straw-colored minnow 116
- sturgeon, smoked 96
- sucker, black 101
- carp 101
- chub 102
- common 101
- hog 101
- spotted 102
- sucker-mouthed minnow 117
- sunfish 87, 88

fishery of the Kaskaskia river—concl'd
 sunfish, long-eared 111
 orange-spotted 110
 survey of Sangamon and Kaskaskia
 fisheries 85
 tackle, kinds of 88
 tadpole cat 118
 tagged fishes 72
 temperatures, air and water 75
 Tom, mad 118
 toothed herring 99
 top minnow 120
 viviparous 121
 trammel-net catches 88
 tributaries 79
 trot-line catches 88
 turtles and frogs 83
Vaillantia camura 120
 value of fishery 83
 vegetation 90
Vigil pellucidus 120
 viviparous top minnow 121
 wall-eyed pike 109
 warmouth bass 111
 water and air temperatures 75
 white crappie 74, 112
 width of Kaskaskia river 74, 77
 Wisconsin glaciation 76, 77
Xenotis megalotis 111
 Yankee Ridge 74, 75
Zygonectes notatus 120
 fishes of Illinois 473-500
 Fox river 476

G

grasshoppers—see Orthoptera
 cultivation to destroy eggs of 269
 poisoning 266
 trapping 268
 Green river 476

H

Hexagenia 90
 hybrid fishes 115, 116, 473, 482, 486-88,
 492-94
 hydroelectric plant 305

I

Illinois, main rivers of 83, 476
 river 476
 State Nurserymen's Association co-
 operates in elm disease study 2
 Illinois-Mississippi canal 476

Iroquois river 476

K

Kankakee river 476
 Kaskaskia island 79, 80
 river 71-123, 476
 Kishwaukee river 476

L

laboratory boat—see Anax
 lakes of Illinois—see fishery, fishes
 Little Wabash river 476
 Lusk creek 476

M

Mackinaw river 476
 Macoupin creek 476
 Middle Fork river 476
 Mississippi river 79, 476

N

new classification—see Plecoptera
 species—see Orthoptera, Plecoptera
 North Fork river 476
 nurseries 61

O

Ohio river 476
 Okaw river—see Kaskaskia river
 Orthoptera 125-279
 collecting localities 136
 new species 230, 237

P

Pecatonica river 476
 Pianezze III-b stain 13
 Plecoptera 281-471
 hydroelectric plant bar rack, nymph
 habitat 305
 new classification 310
 genus 417
 species 365, 423, 441, 445, 447, 455,
 459
 variety 363

R

rivers of Illinois—see by name
 principal 83, 476
 Rock river 476

S

Saline river 476
 Salt creek 476
 Salt Fork river 476
 Sangamon river 476
Saperda tridentata, thought cause of elm
 disease 3

Sargasso sea, spawning ground of eel	V
100	Vermilion river 476
Shoal creek 476	W
Skillet Fork river 476	Wabash river 476
Spoon river 476	Walsh, Benjamin Dann, first state en-
stain, Pianezze III-b 13	tomologist 281
Stoneflies—see Plecoptera	Wilt—see elm disease

ADDITIONAL CURRENT PUBLICATIONS of the Illinois State Natural History Survey

ILLINOIS NATURAL HISTORY SURVEY CIRCULAR

- 24.—Care of Trees. By L. E. Sawyer. February 1932. 7 pp., 5 figs. Contents: Watering, Fertilization, Trimming, Surgery. Fifteen cents.
- 25.—How to Collect and Preserve Insects. By H. H. Ross. February 1934. 27 pp., 11 figs. Partial contents: Where to collect, What to use, How to handle unmounted specimens, How to mount and preserve the collection, How to label the specimens, Housing the collection permanently. Fifteen cents.
- 26.—Insect Enemies of Peach in Illinois. By S. C. Chandler and W. P. Flint. May 1935. 38 pp., 30 figs. Contents: Appearance and type of injury, control, and life history of San Jose scale, peach borer, lesser peach borer, shothole borer and peach tree bark beetle, terrapin scale, oriental fruit moth, peach twig borer, plum curculio, tarnished plant bug, stink bugs, Japanese beetle, green June beetle, cotton leaf worm moth, aphids, grasshoppers, tree crickets, thrips. Twenty-five cents.

ILLINOIS NATURAL HISTORY SURVEY BOOKLIST

Available publications of the ILLINOIS STATE NATURAL HISTORY SURVEY.
March 1935. 13 pp., 257 titles.

Address orders and correspondence to the Chief,
ILLINOIS STATE NATURAL HISTORY SURVEY
Natural History Bldg., Urbana, Ill.

CURRENT PUBLICATIONS

of the Illinois State Natural History Survey

A.—ILLINOIS NATURAL HISTORY SURVEY BULLETIN. Volume XX.

Article I.—Initial Studies of American Elm Diseases in Illinois. By Hubert A. Harris. October 1932. 70 pp., 35 figs. Partial contents: Character of the present elm malady, Laboratory methods, General results of cultural tests, Specific diseases, Other infections, Inoculation trials, Preliminary control experiments. Fifty cents.

Article II.—A Survey of the Fishery of the Kaskaskia River. By Wilbur M. Luce. August 1933. 53 pp., 12 figs. Partial contents: Hydrography, Methods and equipment, Value of the fishery, Kinds and numbers of mussels, Kinds of fishes caught with various tackle, Factors affecting abundance, Rate of growth, Large fishes, Small fishes. Fifty cents.

Article III.—The Dermaptera and Orthoptera of Illinois. By Morgan Hebard. November 1934. 155 pp., front. + 167 figs. Partial contents: Biology and habits of the orders, Ecological factors affecting Orthoptera, Key to the orders, Description of species, Control measures. \$1.

Article IV.—The Stoneflies, or Plecoptera, of Illinois. By Theodore H. Frison. January 1935. 191 pp., front. + 344 figs. Contents: Status of the group; Stonefly structure; Biology of the order; Collecting, rearing and preserving stoneflies; Catalog of species, including classification, description, habits and records of Illinois stoneflies, and new classification of North American forms. \$1.25.

B.—ILLINOIS NATURAL HISTORY SURVEY CIRCULAR.

24.—Care of Trees. By L. E. Sawyer. February 1932. 7 pp., 5 figs. Contents: Watering, Fertilization, Trimming, Surgery. Fifteen cents.

25.—How to Collect and Preserve Insects. By H. H. Ross. February 1934. 27 pp., 11 figs. Partial contents: Where to collect, What to use, How to handle unmounted specimens, How to mount and preserve the collection, How to label the specimens, Housing the collection permanently. Fifteen cents.

26.—Insect Enemies of Peach in Illinois. By S. C. Chandler and W. P. Flint. May 1935. 38 pp., 30 figs. Contents: Appearance and type of injury, control, and life history of San Jose scale, peach borer, lesser peach borer, shothole borer and peach tree bark beetle, terrapin scale, oriental fruit moth, peach twig borer, plum curculio, tarnished plant bug, stink bugs, Japanese beetle, green June beetle, cotton leaf worm moth, aphids, grasshoppers, tree crickets, thrips. Twenty-five cents.

C.—ILLINOIS NATURAL HISTORY SURVEY BOOKLIST.

Available publications of the ILLINOIS STATE NATURAL HISTORY SURVEY.
March 1935. 13 pp., 257 titles.

Address orders and correspondence to the Chief,
ILLINOIS STATE NATURAL HISTORY SURVEY
Natural History Bldg., Urbana, Ill.